arXiv:1109.5328v2 [math.AP] 18 Apr 2012

Global symmetric classical and strong solutions of the full compressible Navier-Stokes equations with vacuum and large initial data

Huanyao Wen, Changjiang Zhu*

School of Mathematics and Statistics
Central China Normal University, Wuhan 430079, China

Abstract

First of all, we get the global existence of classical and strong solutions of the full compressible Navier-Stokes equations in three space dimensions with initial data which is large and spherically or cylindrically symmetric. The appearance of vacuum is allowed. In particular, if the initial data is spherically symmetric, the space dimension can be taken not less than two. The analysis is based on some delicate *a priori* estimates globally in time which depend on the assumption $\kappa = O(1+\theta^q)$ where $q > r$ (r can be zero), which relaxes the condition $q \geq 2+2r$ in [14, 29, 42]. This could be viewed as an extensive work of [18] where the equations hold in the sense of distributions in the set where the density is positive with initial data which is large, discontinuous, and spherically or cylindrically symmetric in three space dimension. Finally, with the assumptions that vacuum may appear and that the solutions are not necessarily symmetric, we establish a blow-up criterion in terms of $\|\rho\|_{L_t^\infty L_x^\infty}$ and $\|\rho\theta\|_{L_t^4 L_x^{\frac{12}{5}}}$ for strong solutions.

Key Words: Compressible Navier-Stokes equations, heat conducting fluids, vacuum, global classical and strong solutions, blow-up criterions.

2000 Mathematics Subject Classification. 35Q30, 35K65, 76N10.

Contents

1 Introduction 2

2 Main results 5

2.1 Global symmetric classical and strong solutions with vacuum 5

2.1.1 Assumptions 5

2.1.2 Global strong solutions 6

2.1.3 Global classical solutions 7

2.2 A blow-up criterion in terms of $\|\rho\|_{L_t^\infty L_x^\infty}$ and $\|\rho\theta\|_{L_t^4 L_x^{\frac{12}{5}}}$ for strong solutions . 7

3 Preliminaries 9

4 Proof of Theorem 2.1.1 10

5 Proof of Theorem 2.1.6 35

---

*Corresponding author. Email: cjzhu@mail.ccnu.edu.cn

6 Proof of Theorem 2.2.2 **43**

References **54**

1 Introduction

The full compressible Navier-Stokes equations can be written in the sense of Eulerian coordinates in $\Omega \subset \mathbb{R}^N$ as follows:

$$\begin{cases} \rho_t + \nabla \cdot (\rho \mathbf{u}) = 0, \\ (\rho \mathbf{u})_t + \operatorname{div}(\rho \mathbf{u} \otimes \mathbf{u}) + \nabla P = \operatorname{div}(\mathcal{T}) + \rho \mathbf{f}, \\ (\rho E)_t + \operatorname{div}(\rho E \mathbf{u}) + \operatorname{div}(P \mathbf{u}) = \operatorname{div}(\mathcal{T} \mathbf{u}) + \operatorname{div}(\kappa \nabla \theta) + \rho \mathbf{u} \cdot \mathbf{f}. \end{cases} \tag{1.1}$$

Here $\mathcal{T}$ is the stress tensor given by

$$\mathcal{T} = \mu \left(\nabla \mathbf{u} + (\nabla \mathbf{u})' \right) + \lambda \operatorname{div} \mathbf{u} I_N,$$

where I_N is a $N \times N$ unit matrix; $\rho = \rho(\mathbf{x}, t)$, $\mathbf{u} = \mathbf{u}(\mathbf{x}, t) = (u_1, \cdots, u_N)(\mathbf{x}, t)$ and $\theta = \theta(\mathbf{x}, t)$ are unknown functions denoting the density, velocity and absolute temperature, respectively; $P = P(\rho, \theta)$, E, $\mathbf{f} = \mathbf{f}(\mathbf{x}, t) = (f_1, \cdots, f_N)(\mathbf{x}, t)$ and κ denote respectively pressure, total energy, external forces and coefficient of heat conduction, where $E = e + \frac{1}{2} u^2$ (e is the internal energy); μ and λ are coefficients of viscosity, satisfying the following physical restrictions:

$$\mu > 0, \ 2\mu + N\lambda \geq 0;$$

P and e satisfy the second principle of thermodynamics:

$$P = \rho^2 \frac{\partial e}{\partial \rho} + \theta \frac{\partial P}{\partial \theta}. \tag{1.2}$$

(1.1) is a well-known model which describes the motion of compressible fluids. There were lots of works on the existence, uniqueness, regularity and asymptotic behavior of the solutions during the last five decades. While, because of the stronger nonlinearity in (1.1) compared with the Navier-Stokes equations for isentropic flow (no temperature equation), many known mathematical results focused on the absence of vacuum (vacuum means $\rho = 0$), refer for instance to [21, 22, 29, 30, 34, 35, 40] for classical solutions. More precisely, the local classical solutions to the Navier-Stokes equations with heat-conducting fluid in Hölder spaces was obtained respectively by Itaya in [21] for Cauchy problem and by Tani in [40] for IBVP with $\inf \rho_0 > 0$, where the space dimension $N = 3$. Using delicate energy methods in Sobolev spaces, Matsumura and Nishida in [34, 35] showed that the global classical solutions exist provided that the initial data is small in some sense and away from vacuum in three space dimension. For large initial data in one space dimension, Kazhikhov, Shelukhi in [30] (for polytropic perfect gas with $\mu, \lambda, \kappa =$const.) and Kawohl in [29] (for real gas with $\kappa = \kappa(\rho, \theta)$, $\mu, \lambda =$ const.) respectively got the global classical solutions to (1.1) in Lagrangian coordinates with $\inf \rho_0 > 0$. The internal energy e and the coefficient of heat conduction κ in [29] satisfy the following assumptions for $\rho \leq \overline{\varrho}$ and $\theta \geq 0$ (we translate these conditions in Eulerian coordinates)

$$\begin{cases} e(\rho, 0) \geq 0, \quad \nu(1 + \theta^r) \leq \partial_\theta e(\rho, \theta) \leq N(\overline{\varrho})(1 + \theta^r), \\ \kappa_0(1 + \theta^q) \leq \kappa(\rho, \theta) \leq \kappa_1(1 + \theta^q), \\ |\partial_\rho \kappa(\rho, \theta)| + |\partial_{\rho\rho} \kappa(\rho, \theta)| \leq \kappa_1(1 + \theta^q), \end{cases} \tag{1.3}$$

where $r \in [0,1]$, $q \geq 2+2r$, and ν, $N(\overline{\varrho})$, κ_0 and κ_1 are positive constants. For the perfect gas (i.e., $P = R\rho\theta$, $e = C_\nu\theta$ for some constants $R > 0$ and $C_\nu > 0$) in the domain exterior to a ball in $\mathbb{R}^N$ ($N = 2$ or 3) with $\mu, \lambda, \kappa =$const., Jiang in [22] got the existence of global spherically symmetric classical large solutions in Hölder spaces.

In fact, Kawohl in [29] also considered the case of density-dependent viscosity for another boundary condition with $\inf \rho_0 > 0$, where $0 < \underline{\mu}_0 \leq \mu(\rho) \leq \overline{\mu}_0$ for any $\rho \geq 0$, and $\underline{\mu}_0$ and $\overline{\mu}_0$ are positive constants. This was generalized to the case $\mu(\rho) = \rho^\alpha$ by Jiang in [23] for $\alpha \in (0, \frac{1}{4})$, and by Qin, Yao in [36] for $\alpha \in (0, \frac{1}{2})$, respectively.

On the existence, asymptotic behavior of the weak solutions of the full compressible Navier-Stokes equations with $\inf \rho_0 > 0$, please refer for instance to [24, 25, 28] for the existence of weak solutions in 1D and for the existence of spherically symmetric weak solutions in $\mathbb{R}^N$ ($N = 2$, 3), and refer to [18] for the existence of spherically and cylindrically symmetric weak solutions in $\mathbb{R}^3$, and refer to [13] for the existence of *variational* solutions in a bounded domain in $\mathbb{R}^N$ ($N = 2$, 3).

In the presence of vacuum (i.e. ρ may vanish), to our best knowledge, the mathematical results on global well-posedness of the full compressible Navier-Stokes equations are usually limited to the existence of weak solutions with special pressure, viscosity and heat conductivity (see [1, 14]). More precisely, Feireisl in [14] got the existence of so-called *variational* solutions in dimension $N \geq 2$. The temperature equation in [14] is satisfied only as an inequality in the sense of distributions. Anyhow, Feireisl's work is the very first attempt towards the existence of weak solutions to the full compressible Navier-Stokes equations in higher dimensions, where the coefficients of viscosity are constants and

$$\begin{cases} \kappa = \kappa(\theta) \in C^2[0,\infty), \ \underline{\kappa}(1+\theta^q) \leq \kappa(\theta) \leq \overline{\kappa}(1+\theta^q) \ \text{ for all } \theta \geq 0, \\ P = P(\rho,\theta) = \mathcal{P}_e(\rho) + \theta\mathcal{P}_\theta(\rho) \ \text{ for all } \rho \geq 0 \text{ and } \theta \geq 0, \\ \mathcal{P}_e, \mathcal{P}_\theta \in C[0,\infty) \cap C^1(0,\infty); \ \mathcal{P}_e(0) = 0, \ \mathcal{P}_\theta(0) = 0, \\ \mathcal{P}_e'(\rho) \geq a_1\rho^{\overline{\gamma}-1} - b_1 \ \text{ for all } \rho > 0; \ \mathcal{P}_e(\rho) \leq a_2\rho^{\overline{\gamma}} + b_1 \ \text{ for all } \rho \geq 0, \\ \mathcal{P}_\theta \ \text{ is non-decreasing in } [0,\infty); \ \mathcal{P}_\theta(\rho) \leq a_3(1+\rho^\Gamma) \ \text{ for all } \rho \geq 0, \end{cases} \tag{1.4}$$

where $\Gamma < \frac{\overline{\gamma}}{2}$ if $N = 2$ and $\Gamma = \frac{\overline{\gamma}}{N}$ for $N \geq 3$; $q \geq 2$, $\overline{\gamma} > 1$, and a_1, a_2, a_3, b_1, $\underline{\kappa}$ and $\overline{\kappa}$ are positive constants. Note that the perfect gas equation of state (i.e. $P = R\rho\theta$ for some constant $R > 0$) is not involved in (1.4). In order that the equations are satisfied as equalities in the sense of distribution, Bresch and Desjardins in [1] proposed some different assumptions from [14], and obtained the existence of global weak solutions to the full compressible Navier-Stokes equations with large initial data in $\mathbb{T}^3$ or $\mathbb{R}^3$. In [1], the viscosity $\mu = \mu(\rho)$ and $\lambda = \lambda(\rho)$ may vanish when vacuum appears, and κ, P and e are assumed to be satisfied

$$\begin{cases} \kappa(\rho,\theta) = \kappa_0(\rho,\theta)(\rho+1)(\theta^q+1), \\ P = R\rho\theta + p_c(\rho), \\ e = C_v\theta + e_c(\rho), \end{cases} \tag{1.5}$$

where $q \geq 2$, R and C_v are two positive constants, $p_c(\rho) = O(\rho^{-\ell})$ and $e_c(\rho) = O(\rho^{-\ell-1})$ (for some $\ell > 1$) when ρ is small enough, and $\kappa_0(\rho,\theta)$ is assumed to satisfy

$$\underline{c}_0 \leq \kappa_0(\rho,\theta) \leq \frac{1}{\underline{c}_0},$$

for $\underline{c}_0 > 0$. On the local existence and uniqueness of strong solutions for $N = 3$, please see [4] for the perfect gas with $\mu, \lambda, \kappa =$const. While, there are no global smooth solutions to (1.1) for Cauchy problem when the initial density is of nontrivially compact support and $\kappa = 0$ (see [44]).

Except for the Cauchy problems with initial density compactly supported, it is still unknown whether the global strong (or classical) solutions exist or not when vacuum appears (i.e., the density may vanish) until recently. In our previous paper [42], we got existence and uniqueness of global classical solutions to the full compressible Navier-Stokes equations in one dimension with large initial data and vacuum. In [42], the coefficient of conduction k depends on the temperature, growing as $1 + \theta^q$ where $q \geq 2 + 2r$ (r can be zero).

As a first step to study the problems (1.1) in high dimensions, we study the problems in high dimensions with some symmetry which reduces the whole system to an one dimensional system with singular source terms. The singularity may be due to $x = 0$, $x = \infty$ or appearance of vacuum, where x is the radius. Our main concern here is to show the existence and uniqueness of global classical and strong solutions to (1.1) with vacuum and initial data which is large, and spherically or cylindrically symmetric in three space dimension. In particular, if the initial data is spherically symmetric, the space dimension can be taken not less than two. This extends the results in [18] where the equations hold in the sense of distributions in the set where the density is positive with initial data which is large, discontinuous, and spherically or cylindrically symmetric in three space dimension. Besides, when the solutions are not necessarily symmetric, we shall establish a blow-up criterion for strong solutions with vacuum.

For compressible isentropic Navier-Stokes equations (i.e. no temperature equation), there are so many results about the well-posedness and asymptotic behaviors of the solutions when vacuum appears. Refer to [15, 27, 31, 33] and [17, 32, 41, 46, 47, 48] for global weak solutions with constant viscosity and with density-dependent viscosity, respectively. Refer to [6, 10] and [2, 3, 5, 37] for global strong solutions and for local strong (classical) solutions with constant viscosity, respectively. Recently, Huang, Li, Xin in [20] and Ding, Wen, Yao, Zhu in [8, 7] independently got existence and uniqueness of global classical solutions, where the initial energy in [20] is assumed to be small in $\mathbb{R}^3$ and $\rho - \widetilde{\rho} \in C\left([0,T]; H^3(\mathbb{R}^3)\right)$, $u \in C\left([0,T]; D^1(\mathbb{R}^3) \cap D^3(\mathbb{R}^3)\right) \cap L^\infty\left([\tau,T]; D^4(\mathbb{R}^3)\right)$ (for $\tau > 0$) which generalized the results in [3], and the initial data in [8, 7] could be large for dimension $N = 1$ and could be large but spherically symmetric for $N \geq 2$, and $(\rho, u) \in C([0,T]; H^4(I))$ (I is bounded in [8], and is bounded or an exterior domain in [7]).

We would like to give some notations which will be used throughout the paper.

Notations:

(i) $I = [a,b]$; $Q_T = I \times [0,T]$ for $T > 0$.

(ii) $\displaystyle\int_\sum f = \int_\sum f\,dx$, for $\sum = I$ or $\mathbb{R}^3$.

(iii) For $1 \leq l \leq \infty$, denote the L^l spaces and the standard Sobolev spaces as follows:

$$L^l = L^l(\Sigma),\ D^{k,l} = \Big\{u \in L^1_{\text{loc}}(\Sigma) : \|\nabla^k u\|_{L^l} < \infty\Big\},$$

$$W^{k,l} = L^l \cap D^{k,l},\ H^k = W^{k,2},\ D^k = D^{k,2},$$

$$D^1_0 = \Big\{u \in L^6 :\ \|\nabla u\|_{L^2} < \infty\},$$

$$\|u\|_{D^{k,l}} = \|\nabla^k u\|_{L^l}.$$

(iv) For two 3×3 matrices $E = (E_{ij}), F = (F_{ij})$, denote the scalar product between E and F by

$$E : F = \sum_{i,j=1}^{3} E_{ij} F_{ij}.$$

(v) $G = (2\mu + \lambda)\mathrm{div} u - P$ is the effective viscous flux.

(vi) $\dot{h} = h_t + u \cdot \nabla h$ denotes the material derivative.

The rest of the paper is organized as follows. In Section 2, we present three main theorems of the paper. In Section 3, some useful lemmas are stated, which will be used to prove the global existence of classical and strong solutions. The main theorems will be proved in Sections 4, 5 and 6, respectively.

2 Main results

2.1 Global symmetric classical and strong solutions with vacuum

Throughout Section 2.1 and the proofs of the main theorems in the section, we take $\Sigma = I$ in the **Notations**. For simplicity, we assume that the external force $\mathbf{f} = 0$. Assume $\Omega = \{\mathbf{x} | a < |\mathbf{x}| < b\}$, for $0 < a < b < \infty$. Then for the symmetric cases, (1.1) takes the form:

$$\begin{cases} \rho_t + (\rho u)_x + \dfrac{m\rho u}{x} = 0, \ \rho \geq 0, \ a < x < b, \ t > 0, \\ \rho u_t + \rho u u_x - \dfrac{\rho v^2}{x} + P_x = \beta(u_{xx} + \dfrac{m u_x}{x} - \dfrac{m u}{x^2}), \\ \rho v_t + \rho u v_x + \dfrac{\rho u v}{x} = \mu(v_{xx} + \dfrac{m v_x}{x} - \dfrac{m v}{x^2}), \\ \rho w_t + \rho u w_x = \mu(w_{xx} + \dfrac{m w_x}{x}), \\ \rho e_t + \rho u e_x + P(u_x + \dfrac{m u}{x}) = (\kappa\theta_x)_x + \dfrac{m\kappa\theta_x}{x} + \wp, \end{cases} \tag{2.1.1}$$

where $\beta = 2\mu + \lambda$, $\wp = \lambda(u_x + \frac{mu}{x})^2 + \mu\left(w_x^2 + 2u_x^2 + (v_x - \frac{mv}{x})^2 + \frac{2mu^2}{x^2}\right)$. In the spherically symmetric case, $m = N - 1$, $x = |\mathbf{x}|$, $\mathbf{u}(\mathbf{x}, t) = u(x, t)\frac{\mathbf{x}}{x}$ and $v = w = 0$. In the cylindrically symmetric case, $m = 1$, $x = \sqrt{x_1^2 + x_2^2}$, and

$$\mathbf{u}(\mathbf{x}, t) = u(x, t)\frac{(x_1, x_2, 0)}{x} + v(x, t)\frac{(-x_2, x_1, 0)}{x} + w(x, t)(0, 0, 1).$$

We consider the initial and boundary conditions:

$$(\rho, \ u, \ v, \ w, \ \theta)\big|_{t=0} = (\rho_0, \ u_0, \ v_0, \ w_0, \ \theta_0)(x) \ \text{ in } \ I, \tag{2.1.2}$$

and

$$(u, v, w, \ \theta_x)\big|_{x=a,b} = 0, \ t \geq 0. \tag{2.1.3}$$

2.1.1 Assumptions

(A_1): $\rho_0 \geq 0$, $\int_I \rho_0 > 0$.

(A_2): μ and λ are constants. $\mu > 0$, $2\mu + (m+1)\lambda > 0$. $e = C_0 Q(\theta) + e_c(\rho)$, $P = \rho Q(\theta) + P_c(\rho)$, $\kappa = \kappa(\theta)$, for some constant $C_0 > 0$. The constant C_0 plays no role in the analysis, we assume

$C_0 = 1$.

(A_3): $P_c(\rho) \geq 0$, $e_c(\rho) \geq 0$, for $\rho \geq 0$; $P_c \in C^2[0,\infty)$; $\rho|\frac{\partial e_c}{\partial \rho}| \leq C_1 e_c(\rho)$, for some constant $C_1 > 0$.

$(A_4) : Q(\cdot) \in C^2[0,\infty)$ satisfies

$$\begin{cases} C_2\left(\beta + (1-\beta)\theta + \theta^{1+r}\right) \leq Q(\theta) \leq C_3\left(\beta + (1-\beta)\theta + \theta^{1+r}\right), \\ C_4(1+\theta^r) \leq Q'(\theta) \leq C_5(1+\theta^r), \end{cases}$$

for some constants $C_i > 0$ ($i = 2, 3, 4, 5$) and $r \geq 0$, $\beta = 0$ or 1.

(A_5): $\kappa \in C^2[0,\infty)$ satisfies

$$C_6(1+\theta^q) \leq \kappa(\theta) \leq C_7(1+\theta^q),$$

for $q > r$, and some constants $C_i > 0$ ($i = 6, 7$).

2.1.2 Global strong solutions

Theorem 2.1.1 *(Strong solutions) In addition to* (A_1)*-*(A_5)*, we assume* $\rho_0 \geq 0$*,* $\rho_0 \in H^2$*,* $u_0 \in H^2 \cap H_0^1$*,* $\theta_0 \in H^2$*,* $\partial_x\theta_0|_{x=0,1} = 0$*, and that the following compatibility conditions are valid:*

$$\begin{cases} \beta(u_{0xx} + \dfrac{mu_{0x}}{x} - \dfrac{mu_0}{x^2}) - P_x(\rho_0,\theta_0) = \sqrt{\rho}_0 g_1, \\ \mu(v_{0xx} + \dfrac{mv_{0x}}{x} - \dfrac{mv_0}{x^2}) = \sqrt{\rho}_0 g_2, \\ \mu(w_{0xx} + \dfrac{mw_{0x}}{x}) = \sqrt{\rho}_0 g_3, \\ (\kappa(\theta_0)\theta_{0x})_x + \dfrac{m\kappa(\theta_0)\theta_{0x}}{x} + \wp(x,0) = \sqrt{\rho_0}\ g_4,\ x \in I, \end{cases} \tag{2.1.4}$$

for some $g_i \in L^2$*,* $i = 1,2,3,4$*. Then there exists a unique global solution* $(\rho,\ u,\ v,\ w,\ \theta)$ *to (2.1.1)-(2.1.3) such that for any* $T > 0$

$$\rho \in C([0,T];H^2),\ (u,v,w,\theta) \in C([0,T];H^2) \cap L^2([0,T];H^3),$$
$$(\sqrt{\rho}u_t, \sqrt{\rho}v_t, \sqrt{\rho}w_t, \sqrt{\rho}e_t) \in L^\infty([0,T];L^2),\ (u_t,v_t,w_t) \in L^2([0,T];H_0^1),\ \theta_t \in L^2([0,T];H^1).$$

Remark 2.1.2 *From the assumptions* (A_2)*-*(A_4)*, we know that the polytropic perfect gas (i.e.,* $P = R\rho\theta$*,* $e = C_\nu\theta$ *for some constants* $R > 0$ *and* $C_\nu > 0$*) is included if we take* $r = \beta = 0$ *and* $e_c(\rho) = P_c(\rho) \equiv 0$ *and* $Q = \frac{C_\nu\theta}{C_0}$*.*

Remark 2.1.3 *The global existence of strong solutions depends on* $q > r$ *in our analysis. For the polytropic perfect gas, Theorem 2.1.1 works for any* $q > 0$*, which relaxes the restriction* $q \geq 2$ *in [14, 29, 42].*

Remark 2.1.4 *Some similar compatibility conditions as (2.1.4) can be referred to [4] and references therein. In [4], the* local H^2-regularity *of* u *and* θ *for* the polytropic perfect gas *was obtained. The detailed reasons why such conditions were needed can be found in [4]. Roughly speaking,* g_1*,* g_2*,* g_3 *and* g_4 *are equivalent to* $\sqrt{\rho}u_t$*,* $\sqrt{\rho}v_t$*,* $\sqrt{\rho}w_t$ *and* $\sqrt{\rho}e_t$ *at* $t = 0$*, respectively.*

Remark 2.1.5 *From the derivation of the Navier-Stokes equations from the Boltzmann equation through the Chapman-Enskog expansion to the second order (see [45] and references therein), we know that* $\mu = \mu(\theta)$*,* $\lambda = \lambda(\theta)$ *and* $\kappa = \kappa(\theta)$*. As in [14, 29, 42],* μ *and* λ *are assumed to be constants here, because of the restrictions of mathematical technique.*

2.1.3 Global classical solutions

Theorem 2.1.6 *(Classical solutions) In addition to* (A_1)*-*(A_5)*, we assume* $\rho_0 \geq 0$*,* $\rho_0 \in H^3$*,* $(\sqrt{\rho_0})_x \in L^\infty$*,* $u_0 \in H^3 \cap H^1_0$*,* $\theta_0 \in H^3$*,* $\partial_x\theta_0|_{x=0,1} = 0$*,* $(Q, P_c, \kappa) \in W^{3,\infty}$*, and that the compatibility conditions (2.1.4) are satisfied for some* $g_i \in L^2$*,* $i = 1, 2, 3, 4$*, and* $\sqrt{\rho_0}g_j \in H^1_0$*,* $j = 1, 2, 3$*, and* $\sqrt{\rho_0}g_4 \in H^1$*. Then there exists a unique global solution* $(\rho,\ u,\ v,\ w,\ \theta)$ *to (2.1.1)-(2.1.3) such that for any* $T > 0$

$$\begin{gathered}
\rho \in C([0,T]; H^3),\ (u, v, w, \theta) \in L^\infty([0,T]; H^3) \cap L^2([0,T]; H^4),\\
(\sqrt{\rho}u_t, \sqrt{\rho}v_t, \sqrt{\rho}w_t, \sqrt{\rho}e_t) \in L^\infty([0,T]; L^2),\ (\rho u_t, \rho v_t, \rho w_t) \in L^\infty([0,T]; H^1_0),\\
(u_t, v_t, w_t) \in L^2([0,T]; H^1_0),\ \theta_t \in L^2([0,T]; H^1),\ ,\rho e_t \in L^\infty([0,T]; H^1).
\end{gathered}$$

Remark 2.1.7 *If* $\rho_0 \in H^4$*, and* $(u_0,\ v_0,\ w_0)$ *satisfies a stronger compatibility conditions*

$$\begin{cases}
\beta(u_{0xx} + \dfrac{mu_{0x}}{x} - \dfrac{mu_0}{x^2}) - P_x(\rho_0, \theta_0) = \rho_0\widetilde{g}_1,\\
\mu(v_{0xx} + \dfrac{mv_{0x}}{x} - \dfrac{mv_0}{x^2}) = \rho_0\widetilde{g}_2,\\
\mu(w_{0xx} + \dfrac{mw_{0x}}{x}) = \rho_0\widetilde{g}_3,\ x \in I,
\end{cases} \tag{2.1.5}$$

for some $\widetilde{g}_i \in H^1_0$ *and* $(\sqrt{\rho_0}\partial_x\widetilde{g}_i)_x \in L^2$*, we can obtain by using the similar arguments as in [42] that* $\rho \in C([0,T]; H^4)$ *and* $(u, v, w) \in C([0,T]; H^4) \cap L^2([0,T]; H^5)$*.*

2.2 A blow-up criterion in terms of $\|\rho\|_{L^\infty_t L^\infty_x}$ and $\|\rho\theta\|_{L^4_t L^{\frac{12}{5}}_x}$ for strong solutions

Throughout Section 2.2 and the proofs of the main theorems in the section, we take $\Sigma = \mathbb{R}^3$ in the **Notations**. In order to establish some sharp blow-up criterions, we only consider that κ =constant, and that the state equations of P and e is of ideal polytropic gas type: $P = a\rho\theta,\ e = C_0\theta$, where a and C_0 are two positive constants. The constants a, C_0 and κ in the equations play no roles in the section, we assume $a = C_0 = \kappa = 1$. If the solutions are regular enough (such as strong solutions), (1.1) is equivalence to the following system:

$$\begin{cases}
\rho_t + \nabla\cdot(\rho u) = 0,\\
\rho u_t + \rho u\cdot\nabla u + \nabla P(\rho,\theta) = \mu\Delta u + (\mu+\lambda)\nabla \mathrm{div} u,\\
\rho\theta_t + \rho u\cdot\nabla\theta + \rho\theta\mathrm{div} u = \frac{\mu}{2}\left|\nabla u + (\nabla u)'\right|^2 + \lambda(\mathrm{div} u)^2 + \Delta\theta,\ \text{in } \mathbb{R}^3.
\end{cases} \tag{2.2.1}$$

System (2.2.1) is supplemented with initial conditions

$$(\rho, u, \theta)|_{t=0} = (\rho_0, u_0, \theta_0),\ x \in \mathbb{R}^3, \tag{2.2.2}$$

with

$$\rho(x,t) \to 0,\ u(x,t) \to 0,\ \theta(x,t) \to 0,\ \text{as } |x| \to \infty,\ \text{for } t \geq 0. \tag{2.2.3}$$

We give the definition of strong solutions to (2.2.1)-(2.2.3) throughout Section 2.2 and the proofs of main theorem in the section.

Definition 2.2.1 *(Strong solution) For* $T > 0$*,* (ρ, u, θ) *is called a strong solution to the compressible Navier-Stokes equations (2.2.1)-(2.2.3) in* $\mathbb{R}^3 \times [0,T]$*, if for some* $q \in (3, 6]$*,*

$$\begin{gathered}
0 \leq \rho \in C([0,T]; W^{1,q} \cap H^1 \cap L^1),\ \rho_t \in C([0,T]; L^2 \cap L^q),\\
(u,\theta) \in C([0,T]; D^2 \cap D^1_0) \cap L^2(0,T; D^{2,q}),\ (u_t, \theta_t) \in L^2(0,T; D^1_0),\\
(\sqrt{\rho}u_t, \sqrt{\rho}\theta_t) \in L^\infty(0,T; L^2),
\end{gathered}$$

and (ρ, u, θ) *satisfies (2.2.1) a.e. in* $\mathbb{R}^3 \times (0,T]$*.*

We present our main theorem, which is on a blow-up criterion for strong solutions to (2.2.1)-(2.2.3), as follows:

Theorem 2.2.2 *Assume* $\rho_0 \geq 0$, $\rho_0 \in H^1 \cap W^{1,q} \cap L^1$, *for some* $q \in (3,6]$, $(u_0, \theta_0) \in D^2 \cap D_0^1$, *and the following compatibility conditions are satisfied:*

$$\begin{cases} \mu \Delta u_0 + (\mu+\lambda)\nabla \mathrm{div} u_0 - \nabla P(\rho_0, \theta_0) = \sqrt{\rho}_0 g_1, \\ \kappa \Delta \theta_0 + \frac{\mu}{2}\left|\nabla u_0 + (\nabla u_0)'\right|^2 + \lambda(\mathrm{div} u_0)^2 = \sqrt{\rho_0} g_2, \ x \in \mathbb{R}^3, \end{cases} \tag{2.2.4}$$

for some $g_i \in L^2$, $i = 1,2$. *Let* (ρ, u, θ) *be a strong solution to (2.2.1)-(2.2.3) in* $\mathbb{R}^3 \times [0,T]$. *If* $0 < T^* < +\infty$ *is the maximum time of existence of the strong solution, then*

$$\lim \sup_{T \nearrow T^*} \left(\|\rho\|_{L^\infty(0,T;L^\infty)} + \|\rho\theta\|_{L^4(0,T;L^{\frac{12}{5}})} \right) = \infty, \tag{2.2.5}$$

provided $3\mu > \lambda$.

Remark 2.2.3 *Under the conditions of Theorem 2.2.2, the local existence of the strong solutions was obtained in [4]. Thus, the assumption* $T^* > 0$ *makes sense.*

Remark 2.2.4 *For the ideal polytropic gas with* $\kappa =$*constant, we noticed recently that Huang and Li in [19] got the global existence of classical and weak solutions to Cauchy problem of (1.1) in* $\mathbb{R}^3$ *with small initial energy and non-vacuum state at infinity.*

Remark 2.2.5 *Before Theorem 2.2.2, there have been several results on the blow-up criterions for strong solutions to (2.2.1), please refer for instance to [11, 12, 39, 43] and references therein. More precisely,*

- *Fan-Jiang-Ou ([11], 3D)*

$$\lim \sup_{t \nearrow T^\star} \left(\|\theta\|_{L^\infty(0,t;L^\infty)} + \|\nabla u\|_{L^1(0,t;L^\infty)} \right) = \infty, \tag{2.2.6}$$

provided $7\mu > \lambda$. *Here the appearance of vacuum is allowed.*

It is well-known that the bound of $\|\nabla u\|_{L^1(0,t;L^\infty)}$ *yields that* $\|\rho\|_{L^\infty(0,t;L^\infty)}$ *is bounded (see (2.2) in [11]), if the initial density is bounded. When* $\|\nabla u\|_{L^1(0,t;L^\infty)}$ *in (2.2.6) is relaxed by the upper bound of the density, the following blow-up criterions were obtained:*

- *Fang-Zi-Zhang ([12], 2D)*

$$\lim \sup_{t \nearrow T^\star} \left(\|\theta\|_{L^\infty(0,t;L^\infty)} + \|\rho\|_{L^\infty(0,t;L^\infty)} \right) = \infty, \tag{2.2.7}$$

where the appearance of vacuum is allowed;

- *Sun-Wang-Zhang ([39], 3D)*

$$\lim \sup_{t \nearrow T^\star} \left(\|\theta\|_{L^\infty(0,t;L^\infty)} + \|\rho\|_{L^\infty(0,t;L^\infty)} + \left\|\frac{1}{\rho}\right\|_{L^\infty(0,t;L^\infty)} \right) = \infty, \tag{2.2.8}$$

provided $7\mu > \lambda$. *Here the appearance of vacuum is not allowed.*

- *Wen-Zhu ([43], 3D)*

$$\lim \sup_{t \nearrow T^\star} \left(\|\theta\|_{L^\infty(0,t;L^\infty)} + \|\rho\|_{L^\infty(0,t;L^\infty)} \right) = \infty, \tag{2.2.9}$$

provided $3\mu > \lambda$. *Here the appearance of vacuum is allowed.*

Remark 2.2.6 *Theorem 2.2.2 is an extension of our former results in [43] (see (2.2.9)). One of the main ingredients is that the estimates of* $\|\sqrt{\rho}\theta\|_{L^\infty(0,T;L^2)}$ *and* $\|\nabla u\|_{L^\infty(0,T;L^2)}$ *are done together, i.e.,*

$$\sup_{0\le t\le T}\int_{\mathbb{R}^3}(\rho|\theta|^2+|\nabla u|^2)\,dx+\int_0^T\int_{\mathbb{R}^3}(|\nabla\theta|^2+\rho|u_t|^2)\,dxdt\le C.$$

In [43], $\|\sqrt{\rho}\theta\|_{L^\infty(0,T;L^2)}$ *and* $\|\sqrt{\rho}u\|_{L^\infty(0,T;L^2)}$ *were done together, which needed the upper bounds of the temperature and the density.*

3 Preliminaries

The lemmas in the section will be useful in the next two sections.

Lemma 3.1 *([42]) Let* $\Omega=[\widetilde{a},\widetilde{b}]$ *be a bounded domain in* $\mathbb{R}$*, and* ρ *be a non-negative function such that*

$$0<M\le\int_\Omega\rho\le K,$$

for constants $M>0$ *and* $K>0$*. Then*

$$\|v\|_{L^\infty(\Omega)}\le\frac{K}{M}\|v_x\|_{L^1(\Omega)}+\frac{1}{M}\left|\int_\Omega\rho v\right|,$$

for any $v\in H^1(\Omega)$*.*

Remark 3.2 *The version of higher dimensions for Lemma 3.1 can be found in [13] or [14].*

Corollary 3.3 *([42]) Consider the same conditions in Lemma 3.1, and in addition assume* $\Omega=I$*, and*

$$\|\rho v\|_{L^1(I)}\le\overline{c}.$$

Then for any $l>0$*, there exists a positive constant* $C=C(M,K,l,\overline{c})$ *such that*

$$\|v^l\|_{L^\infty(I)}\le C\|(v^l)_x\|_{L^1(I)}+C,$$

for any $v^l\in H^1(I)$*.*

Lemma 3.4 *(Poincaré inequality) For any* $v\in H_0^1(I)$*, we have*

$$\|v\|_{L^\infty(I)}\le\|v_x\|_{L^1}.$$

Lemma 3.5 *([38]). Assume* $X\subset E\subset Y$ *are Banach spaces and* $X\hookrightarrow\hookrightarrow E$*. Then the following imbedding are compact:*

$$(i)\quad\left\{\varphi:\varphi\in L^q(0,T;X),\frac{\partial\varphi}{\partial t}\in L^1(0,T;Y)\right\}\hookrightarrow\hookrightarrow L^q(0,T;E),\quad\text{if}\quad 1\le q\le\infty;$$

$$(ii)\quad\left\{\varphi:\varphi\in L^\infty(0,T;X),\frac{\partial\varphi}{\partial t}\in L^r(0,T;Y)\right\}\hookrightarrow\hookrightarrow C([0,T];E),\quad\text{if}\quad 1<r\le\infty.$$

4 Proof of Theorem 2.1.1

In the section, we denote by C a generic constant depending only on $\|(\rho_0, u_0, v_0, w_0, \theta_0)\|_{H^2}$, $\|g_i\|_{L^2}$ $(i = 1, 2, 3, 4)$, T, λ, μ, a, b, and some other known constants, but independent of the solutions and the lower bounds of the density. We denote by

$$A \lesssim B$$

if there exists a generic constant C such that $A \leq CB$.

The strategies on proving Theorem 2.1.1 are very classical. More precisely, we derive various *a priori* estimates for strong solutions of the Navier-Stokes equations (2.1.1)-(2.1.3), which are independent of positive lower bounds of the initial density. Then we shall construct a sequence of approximate initial data where the initial density has a lower bound $\varepsilon > 0$. With these *a priori* estimates uniform for ε, we take the limits $\varepsilon \to 0^+$.

From now on, for any $T > 0$, we shall derive some delicate *a priori* estimates for the strong solutions (ρ, u, v, w, θ) as in Theorem 2.1.1 with $\inf\limits_{(x,t)\in Q_T} \rho > 0$. These energy estimates will be finished by five steps.

Step 1: Basic energy inequality

Lemma 4.1 *Under the conditions of Theorem 2.1.1, we have for any* $t \in [0, T]$

$$\int_I x^m \rho(1 + e + u^2 + v^2 + w^2) \leq C.$$

Proof. This bound is standard and follow directly from the equations $(1.1)_1$, $(1.1)_3$ and the boundary conditions. □

Step 2: Upper bound of density

Lemma 4.2 *Under the conditions of Theorem 2.1.1, we have*

$$\|\rho\|_{L^\infty(Q_T)} \leq C.$$

Proof. The idea of the proof is essentially that of a similar result of Frid, Shelukhin [16] where $m = 1$, but with a slightly modification. We omit it for brevity. □

Step 3: H^1-estimates of (ρ, u, v, w)

The next lemma plays an important role in the paper, whose proofs are improved in contrast with [42]. The condition $q > r$ instead of $q \geq 2 + 2r$ is enough here.

Lemma 4.3 *Under the conditions of Theorem 2.1.1, for* $q > r$*, and for any* $0 < \alpha < \min\{1, q - r\}$*, assume* $2\mu + (m + 1)\lambda > 0$*, we have*

$$\int_{Q_T} x^m \frac{(1 + \theta^q)\theta_x^2}{\theta^{1+\alpha}} \leq C,$$

where the generic constant C depends on α.

Remark 4.4 *The proofs of this lemma depend on the boundary condition* $\theta_x|_{x=a,b} = 0$.

Proof. From (1.2) and (2.1.1), we get

$$\rho e_\theta \theta_t + \rho u e_\theta \theta_x + \theta P_\theta (u_x + \frac{mu}{x}) = (\kappa(\theta)\theta_x)_x + \frac{m\kappa\theta_x}{x} + \wp. \tag{4.1}$$

Substituting $e = Q(\theta) + e_c(\rho)$ and $P = \rho Q(\theta) + P_c(\rho)$ into (4.1), we get

$$\rho Q'(\theta)\theta_t + \rho u Q'(\theta)\theta_x + \rho\theta Q'(\theta)(u_x + \frac{mu}{x}) = (\kappa(\theta)\theta_x)_x + \frac{m\kappa\theta_x}{x} + \wp, \tag{4.2}$$

or

$$(\rho Q)_t + (\rho u Q)_x + \frac{m\rho u Q}{x} + \rho\theta Q'(\theta)(u_x + \frac{mu}{x}) = (\kappa(\theta)\theta_x)_x + \frac{m\kappa\theta_x}{x} + \wp. \tag{4.3}$$

Multiplying (4.2) by $x^m\theta^{-\alpha}$, and integrating by parts over Q_T, we have

$$\begin{aligned}
&\int_{Q_T} x^m \left(\frac{\alpha\kappa(\theta)\theta_x^2}{\theta^{1+\alpha}} + \frac{\lambda(u_x + \frac{mu}{x})^2}{\theta^\alpha} + \frac{\mu[w_x^2 + 2u_x^2 + (v_x - \frac{mv}{x})^2 + \frac{2mu^2}{x^2}]}{\theta^\alpha} \right) \\
&= \int_I x^m \rho \int_0^\theta \frac{Q'(\xi)}{\xi^\alpha} - \int_I x^m \rho_0 \int_0^{\theta_0} \frac{Q'(\xi)}{\xi^\alpha} + \int_{Q_T} x^m \rho\theta^{1-\alpha} Q'(\theta)(u_x + \frac{mu}{x}) \\
&\lesssim \int_I x^m \rho \left| \int_0^\theta \frac{1+\xi^r}{\xi^\alpha} \right| + \int_I x^m \rho_0 \left| \int_0^{\theta_0} \frac{1+\xi^r}{\xi^\alpha} \right| + \int_{Q_T} x^m \rho\theta^{1-\alpha}(1+\theta^r)|u_x + \frac{mu}{x}| \\
&\lesssim \int_I x^m \rho (1 + \theta^{1+r}) + \int_I x^m \rho_0 (1 + \theta_0^{1+r}) + \int_{Q_T} x^m \rho\theta^{1-\alpha}(1+\theta^r)|u_x + \frac{mu}{x}|,
\end{aligned} \tag{4.4}$$

where we have used (A_4) and Young inequality. Since $\mu > 0$, we have from (4.4), (A_2), (A_4) and Lemma 4.1

$$\begin{aligned}
\alpha \int_{Q_T} x^m \frac{\kappa\theta_x^2}{\theta^{1+\alpha}} \le & C - \int_{Q_T} x^m \frac{\lambda(u_x + \frac{mu}{x})^2 + 2\mu(u_x^2 + \frac{mu^2}{x^2})}{\theta^\alpha} \\
& + C \int_{Q_T} x^m \rho\theta^{1-\alpha}(1+\theta^r)|u_x + \frac{mu}{x}| = I_1 + I_2 + I_3.
\end{aligned} \tag{4.5}$$

Without loss of generality, we assume $\lambda < 0$. In fact, if $\lambda \ge 0$, I_2 is obviously a good term:

$$I_2 \le -2\mu \int_{Q_T} x^m \frac{u_x^2 + \frac{mu^2}{x^2}}{\theta^\alpha}.$$

For $\lambda < 0$, we use Cauchy inequality to get

$$\begin{aligned}
I_2 = & - \int_{Q_T} x^m \frac{(2\mu + \lambda)u_x^2 + (2\mu + m\lambda)\frac{mu^2}{x^2} + \frac{2m\lambda u u_x}{x}}{\theta^\alpha} \\
\le & - [2\mu + (m+1)\lambda] \int_{Q_T} x^m \frac{u_x^2 + \frac{mu^2}{x^2}}{\theta^\alpha}.
\end{aligned} \tag{4.6}$$

For I_3, using Cauchy inequality again, we get

$$\begin{aligned}
I_3 \le & [2\mu + (m+1)\lambda] \int_{Q_T} x^m \frac{u_x^2 + \frac{mu^2}{x^2}}{\theta^\alpha} + C \int_{Q_T} x^m \rho^2 (1 + \theta^{2+2r-\alpha}) \\
\le & [2\mu + (m+1)\lambda] \int_{Q_T} x^m \frac{u_x^2 + \frac{mu^2}{x^2}}{\theta^\alpha} + C \int_0^T \|\theta\|_{L^\infty}^{1+r-\alpha} \int_I x^m \rho\theta^{1+r} + C \\
\le & [2\mu + (m+1)\lambda] \int_{Q_T} x^m \frac{u_x^2 + \frac{mu^2}{x^2}}{\theta^\alpha} + C \int_0^T \|\theta\|_{L^\infty}^{1+r-\alpha} + C.
\end{aligned} \tag{4.7}$$

Substituting (4.6) and (4.7) into (4.5), we have

$$\alpha\int_{Q_T} x^m\frac{\kappa\theta_x^2}{\theta^{1+\alpha}} \le C\int_0^T \|\theta\|_{L^\infty}^{1+r-\alpha}+C. \tag{4.8}$$

Now we estimate the first term of the right hand side of (4.8).

Case 1: $r<q<1+2r-\alpha$

$$\begin{aligned}C\int_0^T \|\theta\|_{L^\infty}^{1+r-\alpha} \lesssim& 1+\int_0^T \|\theta^{r-\alpha}\theta_x\|_{L^2}\\ \le& C+C\int_0^T\left(\int_I \frac{\theta_x^2\theta^q}{\theta^{1+\alpha}}\theta^{2r-\alpha+1-q}\right)^{\frac{1}{2}}\\ \le&\frac{1}{4}\alpha\int_{Q_T} x^m\frac{\kappa\theta_x^2}{\theta^{1+\alpha}}+C\int_0^T\|\theta\|_{L^\infty}^{2r-\alpha+1-q}+C\\ \le&\frac{1}{4}\alpha\int_{Q_T} x^m\frac{\kappa\theta_x^2}{\theta^{1+\alpha}}+\frac{1}{2}C\int_0^T\|\theta\|_{L^\infty}^{1+r-\alpha}+C,\end{aligned}$$

where we have used Corollary 3.3, (A_5) and Young inequality.

This gives

$$C\int_0^T \|\theta\|_{L^\infty}^{1+r-\alpha} \le \frac{1}{2}\alpha\int_{Q_T} x^m\frac{\kappa\theta_x^2}{\theta^{1+\alpha}}+C. \tag{4.9}$$

Case 2: $q\ge 1+2r-\alpha$

Using Young inequality, we have

$$\begin{aligned}C\int_0^T \|\theta\|_{L^\infty}^{1+r-\alpha} \lesssim& 1+\int_0^T \|\theta^{r-\alpha}\theta_x\|_{L^2}\\ \lesssim& 1+\int_0^T\left(\int_I \frac{\theta_x^2\theta^{1+2r-\alpha}}{\theta^{1+\alpha}}\right)^{\frac{1}{2}}\\ \le& C+C\int_0^T\left(\int_I \frac{\kappa\theta_x^2}{\theta^{1+\alpha}}\right)^{\frac{1}{2}}\\ \le&\frac{1}{2}\alpha\int_{Q_T} x^m\frac{\kappa\theta_x^2}{\theta^{1+\alpha}}+C.\end{aligned} \tag{4.10}$$

Substituting (4.9) or (4.10) into (4.8), we complete the proof of Lemma 4.3. □

The next estimate is a corollary of Lemma 4.3, whose proof can be found in [42] (Corollary 3.1). For completeness, we present the proof.

Corollary 4.5 *Under the conditions of Theorem 2.1.1, we have*

$$\int_0^T \|\theta\|_{L^\infty}^{q-\alpha+1}\le C.$$

Proof. By Corollary 3.3, (A_5) and Lemma 4.3, we have

$$\begin{aligned}\int_0^T \|\theta\|_{L^\infty}^{q-\alpha+1} =& \int_0^T \|\theta^{\frac{q-\alpha+1}{2}}\|_{L^\infty}^2\\ \le& C\int_0^T\int_I\left(\theta^{\frac{q-\alpha-1}{2}}\theta_x\right)^2+C\\ =& C\int_0^T\int_I\frac{\theta^q\theta_x^2}{\theta^{\alpha+1}}+C\\ \le& C.\end{aligned}$$

□

Lemma 4.6 *Under the conditions of Theorem 2.1.1, we have*

$$\int_{Q_T} x^m(u_x^2 + v_x^2 + x^{-2}u^2 + x^{-2}v^2) \le C.$$

Proof. Multiplying $(2.1.1)_2$ and $(2.1.1)_3$ by $x^m u$ and $x^m v$ respectively, and integrating by parts over I, we have

$$\frac{1}{2}\frac{d}{dt}\int_I x^m \rho u^2 - \int_I x^{m-1}\rho u v^2 + \int_I x^m u P_x + \beta \int_I x^m(u_x^2 + m x^{-2}u^2) = 0. \tag{4.11}$$

$$\frac{1}{2}\frac{d}{dt}\int_I x^m \rho v^2 + \int_I x^{m-1}\rho u v^2 + \mu \int_I x^m(v_x^2 + m x^{-2}v^2) = 0. \tag{4.12}$$

Adding (4.12) into (4.11), we have

$$\frac{1}{2}\frac{d}{dt}\int_I x^m \rho(u^2+v^2) + \int_I x^m u P_x + \beta \int_I x^m(u_x^2 + m x^{-2}u^2) + \mu \int_I x^m(v_x^2 + m x^{-2}v^2) = 0. \tag{4.13}$$

Integrating (4.13) over $(0,T)$, and using integration by parts and Cauchy inequality, we have

$$\begin{aligned}
&\beta \int_{Q_T} x^m(u_x^2 + m x^{-2}u^2) + \mu \int_{Q_T} x^m(v_x^2 + m x^{-2}v^2)\\
\le&\frac{1}{2}\int_I x^m \rho_0(u_0^2 + v_0^2) + \int_0^T \int_I x^m u_x P + m\int_0^T\int_I x^{m-1}uP\\
\le&C + \int_0^T \int_I x^m(u_x + m x^{-1}u)P\\
\le&\frac{1}{2}\beta \int_{Q_T} x^m(u_x^2 + m x^{-2}u^2) + C\int_{Q_T} x^m P^2.
\end{aligned}$$

This gives

$$\begin{aligned}
&\frac{1}{2}\beta \int_{Q_T} x^m(u_x^2 + m x^{-2}u^2) + \mu \int_{Q_T} x^m(v_x^2 + m x^{-2}v^2)\\
\lesssim&\int_{Q_T} x^m \rho^2(1+\theta^{2+2r}) + 1\\
\lesssim&\int_0^T \|\theta\|_{L^\infty}^{1+r}\int_I x^m \rho \theta^{1+r} + 1\\
\lesssim&\int_0^T \|\theta\|_{L^\infty}^{1+r} + 1,
\end{aligned}$$

where we have used (A_2), (A_3), (A_4) and Lemmas 4.1-4.2.

Since $q > r$ and $0 < \alpha < q - r$, we have $1 + r < q - \alpha + 1$. Thus, using Young inequality and Corollary 4.5, we have

$$\begin{aligned}
&\frac{1}{2}\beta \int_{Q_T} x^m(u_x^2 + m x^{-2}u^2) + \mu \int_{Q_T} x^m(v_x^2 + m x^{-2}v^2)\\
\lesssim&\int_0^T \|\theta\|_{L^\infty}^{q-\alpha+1} + 1 \le C.
\end{aligned}$$

□

Lemma 4.7 *Under the conditions of Theorem 2.1.1, we have for any* $t \in [0,T]$

$$\int_I x^m(v_x^2 + x^{-2}v^2) + \int_{Q_T} x^m \rho v_t^2 \leq C.$$

Proof. Multiplying $(2.1.1)_3$ by $x^m v_t$, integrating by parts over I, and using Cauchy inequality, we have

$$\begin{aligned}
&\frac{\mu}{2}\frac{d}{dt}\int_I x^m(v_x^2 + mx^{-2}v^2) + \int_I x^m \rho v_t^2 \\
=&-\int_I x^m \rho u v_x v_t - \int_I x^{m-1}\rho u v v_t \\
\leq&\frac{1}{2}\int_I x^m \rho v_t^2 + C\int_I x^m \rho u^2 v_x^2 + C\int_I x^{m-2}\rho u^2 v^2.
\end{aligned}$$

Thus, we have

$$\begin{aligned}
&\mu\frac{d}{dt}\int_I x^m(v_x^2 + mx^{-2}v^2) + \int_I x^m \rho v_t^2 \\
\lesssim&\|u\|_{L^\infty}^2\|\rho\|_{L^\infty}\int_I x^m v_x^2 + \|v\|_{L^\infty}^2\int_I x^m \rho u^2 \\
\lesssim&\int_I x^m(u_x^2 + mx^{-2}u^2)\int_I x^m v_x^2 + \int_I x^m(v_x^2 + mx^{-2}v^2),
\end{aligned}$$

where we have used Lemmas 4.1-4.2 and Poincaré inequality.

By Gronwall inequality and Lemma 4.6, we complete the proof of Lemma 4.7. □

Corollary 4.8 *Under the conditions of Theorem 2.1.1, we have*

$$\|v\|_{L^\infty(Q_T)} + \int_{Q_T} x^m v_{xx}^2 \leq C.$$

Proof. This is an immediately result from Lemmas 4.2, 4.6, 4.7, Poincaré inequality and $(2.1.1)_3$. □

Lemma 4.9 *Under the conditions of Theorem 2.1.1, we have for any* $t \in [0,T]$

$$\int_I x^m w_x^2 + \int_{Q_T} x^m \rho w_t^2 \leq C.$$

Proof. Multiplying $(2.1.1)_4$ by $x^m w_t$, integrating by parts over I, and using Cauchy inequality, we have

$$\begin{aligned}
\int_I x^m \rho w_t^2 + \frac{\mu}{2}\frac{d}{dt}\int_I x^m w_x^2 =& -\int_I x^m \rho u w_x w_t \\
\leq&\frac{1}{2}\int_I x^m \rho w_t^2 + \frac{1}{2}\int_I x^m \rho u^2 w_x^2.
\end{aligned}$$

Thus, we apply Lemma 4.2 and Poincaré inequality to get

$$\begin{aligned}
\int_I x^m \rho w_t^2 + \mu\frac{d}{dt}\int_I x^m w_x^2 \leq&\|\rho\|_{L^\infty}\|u\|_{L^\infty}^2\int_I x^m w_x^2 \\
\lesssim&\int_I x^m(u_x^2 + mx^{-2}u^2)\int_I x^m w_x^2.
\end{aligned}$$

Using Gronwall inequality and Lemma 4.6, we complete the proof of Lemma 4.9. □

Similar to Corollary 4.8, we get the next corollary.

Corollary 4.10 *Under the conditions of Theorem 2.1.1, we have*

$$\int_{Q_T} x^m w_{xx}^2 \le C.$$

Lemma 4.11 *Under the conditions of Theorem 2.1.1, we have for any $t \in [0,T]$*

$$\int_I x^m(\rho\theta^{q+r+2} + u_x^2 + x^{-2}u^2) + \int_{Q_T} x^m \left(\rho u_t^2 + (1+\theta^q)^2\theta_x^2\right) \le C.$$

Proof. Multiplying $(2.1.1)_2$ by $x^m u_t$, integrating by parts over I, and using Cauchy inequality, we have

$$\begin{aligned}
&\int_I x^m \rho u_t^2 + \frac{\beta}{2}\frac{d}{dt}\int_I x^m(u_x^2 + mx^{-2}u^2)\\
=&-\int_I x^m \rho u u_x u_t + \int_I x^{m-1}\rho v^2 u_t - \int_I x^m P_x u_t\\
\le&\frac{1}{4}\int_I x^m \rho u_t^2 + C\int_I x^m \rho(u^2u_x^2 + x^{-2}v^4) + \int_I x^m P u_{xt} + m\int_I x^{m-1} P u_t.
\end{aligned}$$

This, along with Lemma 4.2 and Poincaré inequality, deduces

$$\begin{aligned}
&\frac{3}{4}\int_I x^m \rho u_t^2 + \frac{\beta}{2}\frac{d}{dt}\int_I x^m(u_x^2 + mx^{-2}u^2)\\
\le&C\|\rho\|_{L^\infty}\|u\|_{L^\infty}^2\int_I x^m u_x^2 + C\int_I x^{m-2}\rho v^4 + \frac{d}{dt}\int_I x^m P u_x - \int_I x^m P_t u_x + m\int_I x^{m-1} P u_t\\
\le&C\left(\int_I x^m(u_x^2 + mx^{-2}u^2)\right)^2 + \frac{d}{dt}\int_I x^m P u_x + C\int_I x^{m-2}\rho v^4 - \int_I x^m P_t u_x + m\int_I x^{m-1} P u_t \qquad (4.14)\\
=&\sum_{i=1}^{5} II_i.
\end{aligned}$$

For II_3, using Lemma 4.1 and Corollary 4.8, we have

$$II_3 \lesssim \|v\|_{L^\infty}^2 \int_I x^m \rho v^2 \le C. \tag{4.15}$$

For II_4, we have

$$\begin{aligned}
II_4 =& -\beta^{-1}\int_I x^m P_t(\beta u_x - P) - \beta^{-1}\int_I x^m P_t P\\
=& -\beta^{-1}\int_I x^m (\rho Q)_t(\beta u_x - P) - \beta^{-1}\int_I x^m P_c(\rho)_t(\beta u_x - P) - \frac{1}{2\beta}\frac{d}{dt}\int_I x^m P^2\\
=&\sum_{i=1}^{3} II_{4,i}.
\end{aligned}$$

For $II_{4,1}$, using (4.3) and $(2.1.1)_2$, we have

$$\begin{aligned}
II_{4,1} =& -\beta^{-1}\int_I (\beta u_x - P)\left((x^m\kappa\theta_x)_x + x^m\wp - (x^m\rho u Q)_x - x^m\rho\theta Q'(\theta)(u_x + \frac{mu}{x})\right)\\
=&\beta^{-1}\int_I x^m(\beta u_x - P)_x(\kappa\theta_x - \rho u Q) + \beta^{-1}\int_I x^m(\beta u_x - P)\rho\theta Q'(\theta)(u_x + \frac{mu}{x})\\
&-\beta^{-1}\int_I x^m(\beta u_x - P)\wp\\
=&\beta^{-1}\int_I x^m(\rho u_t + \rho u u_x - \frac{\rho v^2}{x} - \frac{m\beta u_x}{x} + \frac{m\beta u}{x^2})(\kappa\theta_x - \rho u Q)\\
&+\beta^{-1}\int_I x^m(\beta u_x - P)\rho\theta Q'(\theta)(u_x + \frac{mu}{x}) - \beta^{-1}\int_I x^m(\beta u_x - P)\wp.
\end{aligned}$$

Recalling $\wp = \lambda(u_x + \frac{mu}{x})^2 + \mu\left(w_x^2 + 2u_x^2 + (v_x - \frac{mv}{x})^2 + \frac{2mu^2}{x^2}\right)$, we have

$$\begin{aligned}
II_{4,1} \le & \frac{1}{8}\int_I x^m \rho u_t^2 + C\int_I x^m \rho u^2 Q^2 + C\int_I x^m (\kappa\theta_x)^2 + C\int_I x^m u^2 u_x^2 + C\int_I x^m \rho v^4 + C\int_I x^m u_x^2 \\
& + C\int_I x^{m-4}u^2 + C\sup_{x\in I}(1+\theta^{1+r})\int_I x^m\left(u_x^2 + \rho^2 Q^2 + \rho + x^{-2}u^2\right) \\
& + C\|\beta u_x - P\|_{L^\infty}\int_I x^m(u_x^2 + x^{-2}u^2 + w_x^2 + v_x^2 + x^{-2}v^2) \\
\le & \frac{1}{8}\int_I x^m \rho u_t^2 + C\int_I x^m(u_x^2 + x^{-2}u^2)\int_I x^m \rho(1+\theta^{2+2r}) + C\int_I x^m(\kappa\theta_x)^2 \\
& + C\left(\int_I x^m(u_x^2 + x^{-2}u^2)\right)^2 + C\sup_{x\in I}(1+\theta^{q-\alpha+1})\int_I x^m\big(u_x^2 + \rho(1+\theta^{2+2r}) + x^{-2}u^2\big) \\
& + C\|\beta u_x - P\|_{L^\infty}\int_I x^m(u_x^2 + x^{-2}u^2 + w_x^2 + v_x^2 + x^{-2}v^2) + C \\
\le & \frac{1}{8}\int_I x^m \rho u_t^2 + C\sup_{x\in I}\theta^{1+r}\int_I x^m(u_x^2 + x^{-2}u^2) + C\int_I x^m(\kappa\theta_x)^2 \\
& + C\left(\int_I x^m(u_x^2 + x^{-2}u^2)\right)^2 + C\sup_{x\in I}(1+\theta^{q-\alpha+1})\int_I x^m\left(u_x^2 + \rho(1+\theta^{q+r+2}) + x^{-2}u^2\right) \\
& + C\|\beta u_x - P\|_{L^\infty}\int_I x^m(u_x^2 + x^{-2}u^2 + w_x^2 + v_x^2 + x^{-2}v^2) + C,
\end{aligned}$$

where we have used Young inequality, Poincaré inequality, Lemmas 4.1-4.2, (A_2), (A_4), (4.15), $q > r$ and $\alpha < q - r$.

This, along with Sobolev inequality, Cauchy inequality, Lemmas 4.2, 4.7, 4.9 and $(2.1.1)_2$, deduces

$$\begin{aligned}
II_{4,1} \le & \frac{1}{8}\int_I x^m \rho u_t^2 + C\int_I x^m(\kappa\theta_x)^2 + C\left(\int_I x^m(u_x^2 + x^{-2}u^2)\right)^2 \\
& + C\sup_{x\in I}(1+\theta^{q-\alpha+1})\left[1 + \int_I x^m\left(u_x^2 + \rho\theta^{q+r+2} + x^{-2}u^2\right)\right] \\
& + C\left(\|(\beta u_x - P)_x\|_{L^2} + \|\beta u_x - P\|_{L^2}\right)\left[1 + \int_I x^m(u_x^2 + x^{-2}u^2)\right] + C \\
\le & \frac{1}{8}\int_I x^m \rho u_t^2 + C\int_I x^m(\kappa\theta_x)^2 + C\left(\int_I x^m(u_x^2 + x^{-2}u^2)\right)^2 \\
& + C\sup_{x\in I}(1+\theta^{q-\alpha+1})\left[1 + \int_I x^m\left(u_x^2 + \rho\theta^{q+r+2} + x^{-2}u^2\right)\right] \\
& + C\left(\|\rho u_t + \rho u u_x - \frac{\rho v^2}{x} - \frac{m\beta u_x}{x} + \frac{m\beta u}{x^2}\|_{L^2} + \|\beta u_x - P\|_{L^2}\right)\left[1 + \int_I x^m(u_x^2 + x^{-2}u^2)\right] + C \\
\le & \frac{1}{4}\int_I x^m \rho u_t^2 + C\int_I x^m(\kappa\theta_x)^2 + C\left(\int_I x^m(u_x^2 + x^{-2}u^2)\right)^2 \\
& + C\sup_{x\in I}(1+\theta^{q-\alpha+1})\left[1 + \int_I x^m\left(u_x^2 + \rho\theta^{q+r+2} + x^{-2}u^2\right)\right] + C.
\end{aligned}$$

For $II_{4,2}$, using $(2.1.1)_1$, integration by parts, Lemma 4.2, (A_3) and (A_4), we have

$$\begin{aligned}
II_{4,2} =&\beta^{-1}\int_I x^m P_c'(\rho)(\rho_x u+\rho u_x+mx^{-1}\rho u)(\beta u_x-P)\\
=&\beta^{-1}\int_I x^m (P_c)_x u(\beta u_x-P)+\beta^{-1}\int_I x^m (P_c)'\rho u_x(\beta u_x-P)+\beta^{-1}\int_I mx^{m-1}(P_c)'\rho u(\beta u_x-P)\\
\le&-\beta^{-1}\int_I x^m P_c u_x(\beta u_x-P)-\beta^{-1}\int_I x^m P_c u(\beta u_x-P)_x-\beta^{-1}\int_I mx^{m-1}P_c u(\beta u_x-P)\\
&+C\int_I x^m(u_x^2+x^{-2}u^2)+C\int_I x^m\rho(1+\theta^{2+2r})+C\\
\le&-\beta^{-1}\int_I x^m P_c u(\beta u_x-P)_x+C\int_I x^m(u_x^2+x^{-2}u^2)+C\int_I x^m\rho(1+\theta^{2+2r})+C.
\end{aligned}$$

Using $(2.1.1)_2$, Lemmas 4.1-4.2, Corollary 4.8, (A_3), Poincaré inequality, Young inequality and $\alpha<q-r$, we have

$$\begin{aligned}
II_{4,2} =&-\beta^{-1}\int_I x^m P_c u(\rho u_t+\rho u u_x-\frac{\rho v^2}{x}-\frac{m\beta u_x}{x}+\frac{m\beta u}{x^2})\\
&+C\int_I x^m(u_x^2+x^{-2}u^2)+C\int_I x^m\rho(1+\theta^{2+2r})+C\\
\le&\frac{1}{4}\int_I x^m\rho u_t^2+C\int_I x^m\rho u^2+C\|u\|_{L^\infty}^2\|\rho\|_{L^\infty}\int_I x^m u_x^2+C\|v\|_{L^\infty}^2\int_I x^m\rho v^2\\
&+C\int_I x^m(u_x^2+x^{-2}u^2)+C\int_I x^m\rho(1+\theta^{2+2r})+C\\
\le&\frac{1}{4}\int_I x^m\rho u_t^2+C\left(\int_I x^m(u_x^2+x^{-2}u^2)\right)^2+C\sup_{x\in I}\theta^{q-\alpha+1}+C.
\end{aligned}$$

Putting all the estimates about $II_{4,1}$ and $II_{4,2}$ into II_4, we have

$$\begin{aligned}
II_4\le&-\frac{1}{2\beta}\frac{d}{dt}\int_I x^m P^2+\frac{1}{2}\int_I x^m\rho u_t^2+C\int_I x^m(\kappa\theta_x)^2+C\left(\int_I x^m(u_x^2+x^{-2}u^2)\right)^2\\
&+C\sup_{x\in I}(1+\theta^{q-\alpha+1})\left[1+\int_I x^m\left(u_x^2+\rho\theta^{q+r+2}+x^{-2}u^2\right)\right]+C.
\end{aligned}\tag{4.16}$$

For II_5, recalling $P=\rho Q+P_c$, we have

$$\begin{aligned}
II_5 =&m\int_I x^{m-1}\rho Q u_t+m\int_I x^{m-1}P_c(\rho)u_t\\
\le&\frac{1}{8}\int_I x^m\rho u_t^2+C\int_I x^m\rho(1+\theta^{2+2r})+C\int_I x^m\rho\\
\le&\frac{1}{8}\int_I x^m\rho u_t^2+C(\|\theta\|_{L^\infty}^{q-\alpha+1}+1)\int_I x^m\rho\theta^{1+r}+C\\
\le&\frac{1}{8}\int_I x^m\rho u_t^2+C\|\theta\|_{L^\infty}^{q-\alpha+1}+C,
\end{aligned}\tag{4.17}$$

where we have used Lemma 4.1, Young inequality and (A_4). Note that $P_c(0)=0$ by (1.2) and (A_2).

Putting (4.15), (4.16) and (4.17) into (4.14), we have

$$\begin{aligned}
&\frac{3}{4}\int_I x^m\rho u_t^2+\frac{\beta}{2}\frac{d}{dt}\int_I x^m(u_x^2+mx^{-2}u^2)\\
\le&C\left(\int_I x^m(u_x^2+mx^{-2}u^2)\right)^2+\frac{d}{dt}\int_I x^mPu_x-\frac{1}{2\beta}\frac{d}{dt}\int_I x^mP^2+\frac{5}{8}\int_I x^m\rho u_t^2+C\int_I x^m(\kappa\theta_x)^2\\
&+C\sup_{x\in I}(1+\theta^{q-\alpha+1})\left[1+\int_I x^m\left(u_x^2+\rho\theta^{q+r+2}+mx^{-2}u^2\right)\right]+C.
\end{aligned}$$

Thus,

$$\begin{aligned}
&\frac{1}{8}\int_I x^m\rho u_t^2+\frac{\beta}{2}\frac{d}{dt}\int_I x^m(u_x^2+mx^{-2}u^2)\\
\le&C\left(\int_I x^m(u_x^2+mx^{-2}u^2)\right)^2+\frac{d}{dt}\int_I x^mPu_x-\frac{1}{2\beta}\frac{d}{dt}\int_I x^mP^2+C\int_I x^m(\kappa\theta_x)^2\\
&+C\sup_{x\in I}(1+\theta^{q-\alpha+1})\left[1+\int_I x^m\left(u_x^2+\rho\theta^{q+r+2}+mx^{-2}u^2\right)\right]+C.
\end{aligned}\tag{4.18}$$

Integrating (4.18) over $(0,t)$ for $t\in[0,T]$, and using Corollary 4.5 and Cauchy inequality, we have

$$\begin{aligned}
&\frac{1}{8}\int_0^t\int_I x^m\rho u_t^2+\frac{\beta}{2}\int_I x^m(u_x^2+mx^{-2}u^2)\\
\le&C\int_0^t\left(\int_I x^m(u_x^2+mx^{-2}u^2)\right)^2+\int_I x^mPu_x+C\int_0^t\int_I x^m(\kappa\theta_x)^2\\
&+C\int_0^t\sup_{x\in I}(1+\theta^{q-\alpha+1})\int_I x^m\left(u_x^2+\rho\theta^{q+r+2}+mx^{-2}u^2\right)+C\\
\le&C\int_0^t\left(\int_I x^m(u_x^2+mx^{-2}u^2)\right)^2+\frac{\beta}{4}\int_I x^mu_x^2+C\int_I x^m(\rho^2Q^2+P_c^2)+C\int_0^t\int_I x^m(\kappa\theta_x)^2\\
&+C\int_0^t\sup_{x\in I}(1+\theta^{q-\alpha+1})\int_I x^m\left(u_x^2+\rho\theta^{q+r+2}+mx^{-2}u^2\right)+C.
\end{aligned}$$

Thus,

$$\begin{aligned}
&\int_0^t\int_I x^m\rho u_t^2+\int_I x^m(u_x^2+mx^{-2}u^2)\\
\le&C\int_0^t\left(\int_I x^m(u_x^2+mx^{-2}u^2)\right)^2+C\int_I x^m\rho\theta^{q+r+2}+C\int_0^t\int_I x^m(\kappa\theta_x)^2\\
&+C\int_0^t\sup_{x\in I}(1+\theta^{q-\alpha+1})\int_I x^m\left(u_x^2+\rho\theta^{q+r+2}+mx^{-2}u^2\right)+C.
\end{aligned}\tag{4.19}$$

The next step is to handle the second term and the third one on the right hand side of (4.19).

Multiplying (4.2) by $x^m \int_0^\theta \kappa(\xi)\, d\xi$, and integrating by parts over I, we have

$$\begin{aligned}
&\frac{d}{dt}\int_I x^m \rho \left(\int_0^\theta Q'(\eta) \int_0^\eta \kappa(\xi) d\xi d\eta \right) + \int_I x^m (\kappa\theta_x)^2 \\
=& -\int_I x^m \rho\theta Q'(\theta)(u_x + mx^{-1}u) \int_0^\theta \kappa(\xi)\, d\xi + \lambda \int_I x^m (u_x + mx^{-1}u)^2 \int_0^\theta \kappa(\xi)\, d\xi \\
&+ \mu \int_I x^m \left(w_x^2 + 2u_x^2 + (v_x - mx^{-1}v)^2 + 2mx^{-2}u^2 \right) \int_0^\theta \kappa(\xi)\, d\xi \qquad (4.20) \\
\lesssim & \|\theta(1+\theta^q)\|_{L^\infty} \left(\int_I x^m \rho (1+\theta^{1+r}) |u_x + mx^{-1}u| + \int_I x^m (w_x^2 + u_x^2 + v_x^2 + x^{-2}v^2 + x^{-2}u^2) \right) \\
\lesssim & \|\theta(1+\theta^q)\|_{L^\infty} \left(1 + (\int_I x^m \rho \theta^{2r+2})^{\frac{1}{2}} \left(\int_I x^m (u_x^2 + x^{-2}u^2) \right)^{\frac{1}{2}} + \int_I x^m (u_x^2 + x^{-2}u^2) \right),
\end{aligned}$$

where we have used (A_4), (A_5), Lemmas 4.1, 4.2, 4.7 and 4.9, Hölder inequality and Cauchy inequality.

By Corollary 3.3, we have

$$\|\theta(1+\theta^q)\|_{L^\infty} \lesssim \|\kappa\theta_x\|_{L^2} + 1. \tag{4.21}$$

Putting (4.21) into (4.20), and using Lemma 4.1 and Cauchy inequality, we have

$$\begin{aligned}
&\frac{d}{dt}\int_I x^m \rho \left(\int_0^\theta Q'(\eta) \int_0^\eta \kappa(\xi) d\xi d\eta \right) + \int_I x^m (\kappa\theta_x)^2 \\
\leq & C(\|\kappa\theta_x\|_{L^2} + 1) \left(1 + (\int_I x^m \rho\theta^{2r+2})^{\frac{1}{2}} \left(\int_I x^m (u_x^2 + x^{-2}u^2) \right)^{\frac{1}{2}} + \int_I x^m (u_x^2 + x^{-2}u^2) \right) (4.22) \\
\leq & \frac{1}{2} \int_I x^m (\kappa\theta_x)^2 + C\|\theta\|_{L^\infty}^{r+1} \int_I x^m (u_x^2 + x^{-2}u^2) + C \left(\int_I x^m (u_x^2 + x^{-2}u^2) \right)^2 + C.
\end{aligned}$$

Integrating (4.22) over $(0, t)$, and using (A_4), (A_5) and Young inequality, we have

$$\begin{aligned}
&\int_I x^m \rho\theta^{q+r+2} + \int_0^t \int_I x^m (\kappa\theta_x)^2 \\
\leq & C \int_0^t \|\theta\|_{L^\infty}^{q-\alpha+1} \int_I x^m (u_x^2 + x^{-2}u^2) + C \int_0^t \left(\int_I x^m (u_x^2 + x^{-2}u^2) \right)^2 + C.
\end{aligned} \tag{4.23}$$

By (4.19), (4.23), Corollary 4.5, Lemma 4.6 and Gronwall inequality, we complete the proof of Lemma 4.11. □

By Poincaré inequality and Lemma 4.11, we get the next corollary.

Corollary 4.12 *Under the conditions of Theorem 2.1.1, we have*

$$\|u\|_{L^\infty(Q_T)} \leq C.$$

Corollary 4.13 *Under the conditions of Theorem 2.1.1, we have*

$$\int_0^T \|\theta\|_{L^\infty}^{2q+2} \leq C.$$

Proof. By Corollary 3.3 and Cauchy inequality, we have

$$\begin{aligned}\int_0^T \sup_{x\in I}\theta^{2q+2} \lesssim & \int_0^T\int_I \theta^{2q+1}|\theta_x| + 1\\ \le & \frac{1}{2}\int_0^T \sup_{x\in I}\theta^{2q+2} + C\int_{Q_T}\theta^{2q}\theta_x^2.\end{aligned}$$

This together with Lemma 4.11 completes the proof of Corollary 4.13. □

Lemma 4.14 *Under the conditions of Theorem 2.1.1, we have for any* $t\in[0,T]$

$$\int_I x^m(\rho_x^2+\rho_t^2) + \int_{Q_T} x^m u_{xx}^2 \le C.$$

Proof. Differentiating $(2.1.1)_1$, we have

$$\rho_{xt} + \rho_{xx}u + 2\rho_x u_x + \rho u_{xx} + mx^{-1}\rho_x u + mx^{-1}\rho u_x - mx^{-2}\rho u = 0. \tag{4.24}$$

Multiplying (4.24) by $2x^m\rho_x$, and integrating by parts over I, we have

$$\begin{aligned}\frac{d}{dt}\int_I x^m\rho_x^2 =& -4\int_I x^m\rho_x^2u_x - \int_I x^m(\rho_x^2)_x u - 2\int_I x^m\rho\rho_x u_{xx}\\ & -2m\int_I x^{m-1}\rho_x^2 u - 2m\int_I x^{m-1}\rho\rho_x u_x + 2m\int_I x^{m-2}\rho\rho_x u\\ =& -3\int_I x^m\rho_x^2u_x - m\int_I x^{m-1}\rho_x^2 u - 2\int_I x^m\rho\rho_x u_{xx}\\ & -2m\int_I x^{m-1}\rho\rho_x u_x + 2m\int_I x^{m-2}\rho\rho_x u\\ =& \sum_{i=1}^5 III_i.\end{aligned} \tag{4.25}$$

For III_1, using Sobolev inequality, Young inequality, (A_2), (A_3), (A_4), and Lemmas 4.1, 4.2, 4.11, we have

$$\begin{aligned}III_1 =& -3\beta^{-1}\int_I x^m\rho_x^2(\beta u_x - P) - 3\beta^{-1}\int_I x^m\rho_x^2 P\\ \lesssim & (\|\beta u_x - P\|_{L^\infty} + \|\rho Q + P_c\|_{L^\infty})\int_I x^m\rho_x^2\\ \lesssim & \left(\|\beta u_x - P\|_{L^2} + \|(\beta u_x - P)_x\|_{L^2} + \|1+\theta^{r+1}\|_{L^\infty}\right)\int_I x^m\rho_x^2\\ \lesssim & \left(1 + \sup_{x\in I}\theta^{q-\alpha+1} + \|(\beta u_x - P)_x\|_{L^2}\right)\int_I x^m\rho_x^2.\end{aligned} \tag{4.26}$$

For $\|(\beta u_x - P)_x\|_{L^2}$, using $(2.1.1)_2$, Lemmas 4.2, 4.11, Corollary 4.8 and Corollary 4.12, we have

$$\begin{aligned}\|(\beta u_x - P)_x\|_{L^2} \lesssim & \|\rho u_t\|_{L^2} + \|\rho u u_x\|_{L^2} + \|x^{-1}\rho v^2\|_{L^2} + \|x^{-1}u_x\|_{L^2} + \|x^{-2}u\|_{L^2}\\ \lesssim & \|\sqrt{\rho}u_t\|_{L^2} + 1.\end{aligned} \tag{4.27}$$

Substituting (4.27) into (4.26), we obtain

$$III_1 \lesssim \left(1 + \sup_{x\in I}\theta^{q-\alpha+1} + \|\sqrt{\rho}u_t\|_{L^2}\right)\int_I x^m\rho_x^2. \tag{4.28}$$

For III_i, i =2, 3, 4, 5, we have

$$III_2 \lesssim \|u\|_{L^\infty} \int_I x^m \rho_x^2 \lesssim \int_I x^m \rho_x^2. \tag{4.29}$$

$$III_3 \lesssim \int_I x^m \rho_x^2 + \int_I x^m u_{xx}^2. \tag{4.30}$$

$$III_4 \lesssim \int_I x^m \rho_x^2 + \int_I x^m u_x^2 \lesssim \int_I x^m \rho_x^2 + 1. \tag{4.31}$$

$$III_5 \lesssim \int_I x^m \rho_x^2 + \int_I x^{m-2} u^2 \lesssim \int_I x^m \rho_x^2 + 1. \tag{4.32}$$

Putting (4.28)-(4.32) into (4.25), we have

$$\frac{d}{dt} \int_I x^m \rho_x^2 \lesssim \left(1 + \sup_{x\in I} \theta^{q-\alpha+1} + \|\sqrt{\rho} u_t\|_{L^2}\right) \int_I x^m \rho_x^2 + \int_I x^m u_{xx}^2 + 1. \tag{4.33}$$

By (4.27), (A_2), (A_3), (A_4) and Lemma 4.2, we have

$$\begin{aligned}\int_I x^m u_{xx}^2 &\lesssim \int_I x^m \rho u_t^2 + \int_I x^m \rho_x^2 Q^2 + \int_I x^m \rho^2 (Q')^2 \theta_x^2 + \int_I x^m \rho_x^2 + 1 \\ &\lesssim \int_I x^m \rho u_t^2 + (1 + \sup_{x\in I} \theta^{2+2r}) \int_I x^m \rho_x^2 + \int_I x^m (1+\theta^{2r}) \theta_x^2 + 1.\end{aligned}$$

Since $q > r$, using Young inequality, we have

$$\int_I x^m u_{xx}^2 \lesssim \int_I x^m \rho u_t^2 + (1 + \sup_{x\in I} \theta^{2+2q}) \int_I x^m \rho_x^2 + \int_I x^m (1+\theta^q)^2 \theta_x^2 + 1. \tag{4.34}$$

Substituting (4.34) into (4.33), and using Corollary 4.5, Lemma 4.11, Corollary 4.13, and Gronwall inequality, we get

$$\int_I x^m \rho_x^2 \le C. \tag{4.35}$$

Substituting (4.35) into (4.34), and using Lemma 4.11 and Corollary 4.13 again, we have

$$\int_{Q_T} x^m u_{xx}^2 \le C.$$

The estimate for ρ_t can be obtained easily by $(2.1.1)_1$. □

Step 4: H^2 estimates of (ρ, u, v, w) and H^1 estimates of θ

Lemma 4.15 *Under the conditions of Theorem 2.1.1, we have for any $t \in [0, T]$*

$$\int_I x^m \rho v_t^2 + \int_{Q_T} x^m \left(v_{xt}^2 + x^{-2} v_t^2\right) \le C.$$

Proof. Differentiating $(2.1.1)_3$ with respect to t, we have

$$\begin{aligned}&\rho v_{tt}+\rho_t v_t+\rho_t u v_x+\rho u_t v_x+\rho u v_{xt}+x^{-1}\rho_t u v+x^{-1}\rho u_t v+x^{-1}\rho u v_t\\ =&\mu(v_{xxt}+mx^{-1}v_{xt}-mx^{-2}v_t).\end{aligned}\tag{4.36}$$

Multiplying (4.36) by $x^m v_t$, and integrating by parts over I, we have

$$\begin{aligned}&\frac{1}{2}\frac{d}{dt}\int_I x^m\rho v_t^2+\mu\int_I x^m(v_{xt}^2+mx^{-2}v_t^2)\\ =&-\frac{1}{2}\int_I x^m\rho_t v_t^2-\int_I x^m(\rho_t u v_x+\frac{\rho_t u v}{x})v_t-\int_I x^m\left(\rho u_t v_x+\rho u v_{xt}+\frac{\rho u_t v+\rho u v_t}{x}\right)v_t\\ =&\sum_{i=1}^{3}IV_i.\end{aligned}\tag{4.37}$$

For IV_1, using $(2.1.1)_1$, integration by parts, Cauchy inequality, Lemma 4.2 and Corollary 4.12, we have

$$\begin{aligned}IV_1=&\frac{1}{2}\int_I(x^m\rho u)_x v_t^2\\ =&-\int_I x^m\rho u v_t v_{xt}\\ \le&\frac{\mu}{6}\int_I x^m v_{xt}^2+C\int_I x^m\rho v_t^2.\end{aligned}\tag{4.38}$$

Similarly, for IV_2, we have

$$\begin{aligned}IV_2=&\int_I(x^m\rho u)_x(uv_x+\frac{uv}{x})v_t\\ =&-\int_I x^m\rho u(u_x v_x+uv_{xx}+\frac{u_x v+uv_x}{x}-\frac{uv}{x^2})v_t-\int_I x^m\rho u(uv_x+\frac{uv}{x})v_{xt}\\ \le&C\int_I x^m\rho v_t^2+C\int_I x^m\rho\left(u^2u_x^2v_x^2+u^4v_{xx}^2+\frac{u^2u_x^2v^2+u^4v_x^2}{x^2}+\frac{u^4v^2}{x^4}\right)\\ &+\frac{\mu}{6}\int_I x^m v_{xt}^2+C\int_I x^m\rho^2u^2(u^2v_x^2+\frac{u^2v^2}{x^2})\\ \le&C\int_I x^m\rho v_t^2+C\|v_x\|_{L^\infty}^2\int_I x^m u_x^2+C\int_I x^m v_{xx}^2+C\|u_x\|_{L^\infty}^2\int_I x^{m-2}v^2\\ &+C\int_I x^m v_x^2+C\int_I x^{m-2}v^2+\frac{\mu}{6}\int_I x^m v_{xt}^2\\ \le&C\int_I x^m\rho v_t^2+C\int_I x^m v_{xx}^2+C\int_I x^m u_{xx}^2+\frac{\mu}{6}\int_I x^m v_{xt}^2+C,\end{aligned}\tag{4.39}$$

where we have used $(2.1.1)_1$, integration by parts, Cauchy inequality, Lemmas 4.2, 4.7, 4.11, Corollary 4.12 and Sobolev inequality.

For IV_3, using Cauchy inequality, Lemmas 4.2, 4.7, Corollary 4.8, Corollary 4.12 and Sobolev inequality, we get

$$\begin{aligned}IV_3=&-\int_I x^m\rho u_t v_x v_t-\int_I x^m\rho u v_{xt}v_t-\int_I x^{m-1}\rho u_t v v_t-\int_I x^{m-1}\rho u v_t^2\\ \le&C\int_I x^m\rho u_t^2+C\int_I x^m\rho v_x^2v_t^2+\frac{\mu}{6}\int_I x^m v_{xt}^2+C\int_I x^m\rho^2u^2v_t^2+C\int_I x^{m-2}\rho v^2v_t^2\\ &+C\int_I x^m\rho v_t^2\\ \le&C\int_I x^m\rho u_t^2+\frac{\mu}{6}\int_I x^m v_{xt}^2+C(1+\int_I x^m v_{xx}^2)\int_I x^m\rho v_t^2.\end{aligned}\tag{4.40}$$

Putting (4.38), (4.39) and (4.40) into (4.37), we have

$$\begin{aligned}
&\frac{d}{dt}\int_I x^m\rho v_t^2+\mu\int_I x^m(v_{xt}^2+mx^{-2}v_t^2)\\
\le& C(1+\int_I x^m v_{xx}^2)\int_I x^m\rho v_t^2+C\int_I x^m v_{xx}^2+C\int_I x^m u_{xx}^2+C\int_I x^m\rho u_t^2+C.
\end{aligned}\tag{4.41}$$

It follows from (4.41), Corollary 4.8, Lemmas 4.11, 4.14, the compatibility conditions and Gronwall inequality, we complete the proof of Lemma 4.15. □

By $(2.1.1)_3$, Lemmas 4.2, 4.7, 4.11, 4.14, 4.15, Corollary 4.8, Corollary 4.12, and Sobolev inequality, we get the next estimate.

Corollary 4.16 *Under the conditions of Theorem 2.1.1, we have for any* $t\in[0,T]$

$$\|v\|_{W^{1,\infty}(Q_T)}+\int_I x^m v_{xx}^2+\int_{Q_T} x^m v_{xxx}^2\le C.$$

Similar to Lemma 4.15 and Corollary 4.16, we obtain the next lemma and the next corollary.

Lemma 4.17 *Under the conditions of Theorem 2.1.1, we have for any* $t\in[0,T]$

$$\int_I x^m\rho w_t^2+\int_{Q_T} x^m w_{xt}^2\le C.$$

Corollary 4.18 *Under the conditions of Theorem 2.1.1, we have for any* $t\in[0,T]$

$$\|w\|_{W^{1,\infty}(Q_T)}+\int_I x^m w_{xx}^2+\int_{Q_T} x^m w_{xxx}^2\le C.$$

Lemma 4.19 *Under the conditions of Theorem 2.1.1, we have for any* $t\in[0,T]$

$$\int_I x^m\left(\rho u_t^2+(1+\theta^q)^2\theta_x^2\right)+\int_{Q_T} x^m\left(u_{xt}^2+x^{-2}u_t^2+\rho(1+\theta^{q+r})\theta_t^2\right)\le C.$$

Proof. Differentiating $(2.1.1)_2$ with respect to t, we have

$$\rho u_{tt}+\rho_t u_t+\rho_t uu_x+\rho u_t u_x+\rho uu_{xt}-\frac{\rho_t v^2}{x}-\frac{2\rho vv_t}{x}+P_{xt}=\beta\left(u_{xxt}+\frac{mu_{xt}}{x}-\frac{mu_t}{x^2}\right).\tag{4.42}$$

Multiplying (4.42) by $x^m u_t$, and integrating by parts over I, we have

$$\begin{aligned}
&\frac{1}{2}\frac{d}{dt}\int_I x^m\rho u_t^2+\beta\int_I x^m(u_{xt}^2+\frac{mu_t^2}{x^2})\\
=&-\frac{1}{2}\int_I x^m\rho_t u_t^2-\int_I x^m\rho_t uu_x u_t+\int_I x^{m-1}\rho_t v^2u_t-\int_I x^m\rho(u_tu_x+uu_{xt}-\frac{2vv_t}{x})u_t\\
&+\int_I P_t(x^m u_{xt}+mx^{m-1}u_t)\\
=&\sum_{i=1}^{5}V_i.
\end{aligned}\tag{4.43}$$

For V_1, using $(2.1.1)_1$, integration by parts, Cauchy inequality, Lemma 4.2, Corollary 4.12, we have

$$\begin{aligned} V_1 =&\frac{1}{2}\int_I (x^m\rho u)_x u_t^2 \\ =&-\int_I x^m \rho u u_t u_{xt} \\ \leq&\frac{\beta}{8}\int_I x^m u_{xt}^2 + C\int_I x^m\rho u_t^2. \end{aligned} \tag{4.44}$$

For V_2 and V_3, using $(2.1.1)_1$, integration by parts and Cauchy inequality again, along with Sobolev inequality, Lemmas 4.2, 4.7, 4.11, Corollary 4.8 and Corollary 4.12, we get

$$\begin{aligned} V_2+V_3 =&\int_I (x^m\rho u)_x (u u_x u_t - \frac{v^2 u_t}{x}) \\ =&-\int_I x^m\rho u(u_x^2 u_t + u u_{xx} u_t + u u_x u_{xt} - \frac{2v v_x u_t + v^2 u_{xt}}{x} + \frac{v^2 u_t}{x^2}) \\ \leq& C\int_I x^m \rho u_t^2 u_x^2 + C\int_I x^m\rho u^2 u_x^2 + C\int_I x^m \rho u_t^2 + C\int_I x^m\rho u^4 u_{xx}^2 + \frac{\beta}{8}\int_I x^m u_{xt}^2 \\ &+ C\int_I x^m\rho^2 u^4 u_x^2 + C\int_I x^m\rho u^2 v^2 v_x^2 + C\int_I x^m \rho^2 u^2 v^4 + C\int_I x^m\rho u^2 v^4 \\ \leq& C(1+\int_I u_{xx}^2)\int_I x^m\rho u_t^2 + C\int_I x^m u_{xx}^2 + \frac{\beta}{8}\int_I x^m u_{xt}^2 + C. \end{aligned} \tag{4.45}$$

For V_4, we have

$$\begin{aligned} V_4 \leq& \|u_x\|_{L^\infty}\int_I x^m\rho u_t^2 + \frac{\beta}{8}\int_I x^m u_{xt}^2 + C\int_I x^m\rho^2 u^2 u_t^2 + C\int_I x^m\rho u_t^2 + C\int_I x^{m-2}\rho v^2 v_t^2 \\ \leq& \frac{\beta}{8}\int_I x^m u_{xt}^2 + C(1+\|u_x\|_{H^1})\int_I x^m\rho u_t^2 + C, \end{aligned} \tag{4.46}$$

where we have used Sobolev inequality, Lemmas 4.2, 4.15, Corollary 4.8 and Corollary 4.12.

For V_5, using Young inequality, (A_2), (A_3), (A_4), and Lemmas 4.2, 4.14, we have

$$\begin{aligned} V_5 \leq& \frac{\beta}{8}\int_I x^m(u_{xt}^2 + \frac{m u_t^2}{x^2}) + C\int_I x^m P_t^2 \\ \leq& \frac{\beta}{8}\int_I x^m(u_{xt}^2 + \frac{m u_t^2}{x^2}) + C\int_I x^m \rho_t^2(\partial_\rho P)^2 + C\int_I x^m\theta_t^2(\partial_\theta P)^2 \\ \leq& \frac{\beta}{8}\int_I x^m(u_{xt}^2 + \frac{m u_t^2}{x^2}) + C(1+\sup_{x\in I}\theta^{2+2r})\int_I x^m\rho_t^2 + C\int_I x^m\rho^2(1+\theta^{2r})\theta_t^2 \\ \leq& \frac{\beta}{8}\int_I x^m(u_{xt}^2 + \frac{m u_t^2}{x^2}) + C\sup_{x\in I}\theta^{2+2q} + C\int_I x^m\rho(1+\theta^{q+r})\theta_t^2 + C. \end{aligned} \tag{4.47}$$

Putting (4.44), (4.45), (4.46) and (4.47) into (4.43), we have

$$\begin{aligned} &\frac{d}{dt}\int_I x^m\rho u_t^2 + \beta\int_I x^m(u_{xt}^2 + \frac{m u_t^2}{x^2}) \\ \leq& C(1+\int_I u_{xx}^2)\int_I x^m\rho u_t^2 + C\int_I x^m u_{xx}^2 + C\sup_{x\in I}\theta^{2+2q} + C\int_I x^m\rho(1+\theta^{q+r})\theta_t^2 + C. \end{aligned} \tag{4.48}$$

Integrating (4.48) over $(0,t)$, and using the compatibility conditions, Lemma 4.14 and Corollary 4.13, we obtain

$$\begin{aligned}&\int_I x^m\rho u_t^2+\int_0^t\int_I x^m(u_{xt}^2+\frac{mu_t^2}{x^2})\\ \le& C\int_0^t(1+\int_I u_{xx}^2)\int_I x^m\rho u_t^2+C\int_0^t\int_I x^m\rho(1+\theta^{q+r})\theta_t^2+C.\end{aligned}\tag{4.49}$$

The next step is to estimate the second integral of the right hand side of (4.49).

Multiplying (4.2) by $x^m\left(\int_0^\theta\kappa(\xi)\,d\xi\right)_t$ (i.e., $x^m\kappa(\theta)\theta_t$), and integrating by parts over I, we have

$$\begin{aligned}&\int_I x^m\rho Q'\kappa\theta_t^2+\frac{1}{2}\frac{d}{dt}\int_I x^m\kappa^2\theta_x^2\\ =&-\int_I x^m\rho uQ'\theta_x\kappa\theta_t-\int_I x^m\rho\theta Q'(u_x+\frac{mu}{x})\kappa\theta_t+\lambda\int_I x^m(u_x+\frac{mu}{x})^2\left(\int_0^\theta\kappa(\xi)\,d\xi\right)_t\\ &+\mu\int_I x^m\left(w_x^2+2u_x^2+(v_x-\frac{mv}{x})^2+\frac{2mu^2}{x^2}\right)\left(\int_0^\theta\kappa(\xi)\,d\xi\right)_t\\ =&\sum_{i=1}^4 VI_i.\end{aligned}\tag{4.50}$$

For VI_1, using Cauchy inequality, (A_4), (A_5), Lemma 4.2 and Corollary 4.12, we have

$$\begin{aligned}VI_1\le&\frac{1}{4}\int_I x^m\rho Q'\kappa\theta_t^2+C\int_I x^m\rho Q'\kappa u^2\theta_x^2\\ \le&\frac{1}{4}\int_I x^m\rho Q'\kappa\theta_t^2+C\int_I x^m(1+\theta^q)^2\theta_x^2.\end{aligned}\tag{4.51}$$

For VI_2, using Cauchy inequality, (A_4) and (A_5) again, along with Lemmas 4.1, 4.11 and Corollary 4.12, we have

$$\begin{aligned}VI_2\le&\frac{1}{4}\int_I x^m\rho Q'\kappa\theta_t^2+C\int_I x^m\rho Q'\kappa\theta^2(u_x^2+\frac{u^2}{x^2})\\ \le&\frac{1}{4}\int_I x^m\rho Q'\kappa\theta_t^2+C(\|u_x\|_{L^\infty}^2+\|u\|_{L^\infty}^2)\int_I x^m\rho(1+\theta^{q+r+2})\\ \le&\frac{1}{4}\int_I x^m\rho Q'\kappa\theta_t^2+C\int_I x^m u_{xx}^2+C.\end{aligned}\tag{4.52}$$

For VI_3, we have

$$\begin{aligned}VI_3\le&\frac{d}{dt}\left(\lambda\int_I x^m(u_x+\frac{mu}{x})^2\int_0^\theta\kappa(\xi)\,d\xi\right)-2\lambda\int_I x^m(u_x+\frac{mu}{x})(u_{xt}+\frac{mu_t}{x})\int_0^\theta\kappa(\xi)\,d\xi\\ \le&\lambda\frac{d}{dt}\left(\int_I x^m(u_x+\frac{mu}{x})^2\int_0^\theta\kappa(\xi)\,d\xi\right)+C\|\theta(1+\theta^q)\|_{L^\infty}\left(\int_I x^m(u_{xt}^2+\frac{mu_t^2}{x^2})\right)^{\frac{1}{2}}\\ \le&\lambda\frac{d}{dt}\left(\int_I x^m(u_x+\frac{mu}{x})^2\int_0^\theta\kappa(\xi)\,d\xi\right)+C(\|\kappa\theta_x\|_{L^2}+1)\left(\int_I x^m(u_{xt}^2+\frac{mu_t^2}{x^2})\right)^{\frac{1}{2}},\end{aligned}\tag{4.53}$$

where we have used Hölder inequality, Lemma 4.11, (4.21) and (A_5).

For VI_4, using Hölder inequality, (4.21) and Lemma 4.11 again, along with Lemmas 4.7, 4.9, we have

$$\begin{aligned}
VI_4 =& \mu\frac{d}{dt}\int_I x^m\left(w_x^2+2u_x^2+(v_x-\frac{mv}{x})^2+\frac{2mu^2}{x^2}\right)\int_0^\theta \kappa(\xi)\,d\xi\\
&-2\mu\int_I x^m\left(w_xw_{xt}+2u_xu_{xt}+(v_x-\frac{mv}{x})(v_{xt}-\frac{mv_t}{x})+\frac{2muu_t}{x^2}\right)\int_0^\theta \kappa(\xi)\,d\xi\\
\le&\mu\frac{d}{dt}\int_I x^m\left(w_x^2+2u_x^2+(v_x-\frac{mv}{x})^2+\frac{2mu^2}{x^2}\right)\int_0^\theta \kappa(\xi)\,d\xi+C(\|\kappa\theta_x\|_{L^2}+1)\left(\int_I x^m w_{xt}^2\right)^{\frac12}\\
&+C(\|\kappa\theta_x\|_{L^2}+1)\left[\left(\int_I x^m u_{xt}^2\right)^{\frac12}+\left(\int_I x^m(v_{xt}^2+\frac{v_t^2}{x^2})\right)^{\frac12}+\left(\int_I x^{m-2}u_t^2\right)^{\frac12}\right].
\end{aligned}\tag{4.54}$$

Substituting (4.51), (4.52), (4.53) and (4.54) into (4.50), we have

$$\begin{aligned}
&\int_I x^m\rho Q'\kappa\theta_t^2+\frac{d}{dt}\int_I x^m\kappa^2\theta_x^2\\
\le&2\mu\frac{d}{dt}\int_I x^m\left(w_x^2+2u_x^2+(v_x-\frac{mv}{x})^2+\frac{2mu^2}{x^2}\right)\int_0^\theta \kappa(\xi)\,d\xi\\
&+2\lambda\frac{d}{dt}\left(\int_I x^m(u_x+\frac{mu}{x})^2\int_0^\theta \kappa(\xi)\,d\xi\right)+C\int_I x^m(1+\theta^q)^2\theta_x^2\\
&+C\int_I x^m u_{xx}^2+C(\|\kappa\theta_x\|_{L^2}+1)\left(\int_I x^m w_{xt}^2\right)^{\frac12}\\
&+C(\|\kappa\theta_x\|_{L^2}+1)\left[\left(\int_I x^m u_{xt}^2\right)^{\frac12}+\left(\int_I x^m(v_{xt}^2+\frac{v_t^2}{x^2})\right)^{\frac12}+\left(\int_I x^{m-2}u_t^2\right)^{\frac12}\right]+C.
\end{aligned}\tag{4.55}$$

Integrating (4.55) over $(0,t)$, and using (A_4), (A_5) and Lemma 4.14, we have

$$\begin{aligned}
&\int_0^t\int_I x^m\rho(1+\theta^{q+r})\theta_t^2+\int_I x^m(1+\theta^q)^2\theta_x^2\\
\lesssim&\int_I x^m\left(w_x^2+u_x^2+v_x^2+\frac{v^2}{x^2}+\frac{u^2}{x^2}\right)\int_0^\theta \kappa(\xi)\,d\xi+\int_0^t\int_I x^m(1+\theta^q)^2\theta_x^2\\
&+\int_0^t(\|\kappa\theta_x\|_{L^2}+1)\left(\int_I x^m w_{xt}^2\right)^{\frac12}+\int_0^t(\|\kappa\theta_x\|_{L^2}+1)\left(\int_I x^m u_{xt}^2\right)^{\frac12}\\
&+\int_0^t(\|\kappa\theta_x\|_{L^2}+1)\left[\left(\int_I x^m(v_{xt}^2+\frac{v_t^2}{x^2})\right)^{\frac12}+\left(\int_I x^{m-2}u_t^2\right)^{\frac12}\right]+1.
\end{aligned}\tag{4.56}$$

Using (4.21), (4.56), Cauchy inequality and Lemmas 4.15, 4.17, we have

$$\begin{aligned}
&\int_0^t\int_I x^m\rho(1+\theta^{q+r})\theta_t^2+\int_I x^m(1+\theta^q)^2\theta_x^2\\
\le&C\int_0^t\int_I x^m(1+\theta^q)^2\theta_x^2+C\int_0^t(\|\kappa\theta_x\|_{L^2}+1)\left[\left(\int_I x^m u_{xt}^2\right)^{\frac12}+\left(\int_I x^{m-2}u_t^2\right)^{\frac12}\right]+C.
\end{aligned}\tag{4.57}$$

By (4.49), (4.57), Cauchy inequality and Gronwall inequality, we complete the proof of Lemma 4.19. □

By Lemmas 3.1 and 4.19, we get the next corollary.

Corollary 4.20 *Under the conditions of Theorem 2.1.1, we have*

$$\|\theta\|_{L^\infty(Q_T)} \le C.$$

Corollary 4.21 *Under the conditions of Theorem 2.1.1, we have for any* $t \in [0,T]$

$$\|u\|_{W^{1,\infty}(Q_T)} + \int_I x^m u_{xx}^2 + \int_{Q_T} x^m \theta_{xx}^2 \le C.$$

Proof. *By (4.34), Lemma 4.14, Lemma 4.19 and Corollary 4.20, we get*

$$\int_I x^m u_{xx}^2 \le C. \tag{4.58}$$

It follows from (4.58), Lemma 4.11, Corollary 4.12 and Sobolev inequality, we obtain

$$\|u\|_{W^{1,\infty}} \le C.$$

By (4.2), (4.58), (A_4), (A_5), Lemmas 4.2, 3.4, 4.11, 4.19, Corollaries 4.12, 4.18, 4.20, and Cauchy inequality, we have

$$\begin{aligned}
\int_I x^m \theta_{xx}^2 \lesssim & \int_I x^m \rho\theta_t^2 + \int_I x^m \theta_x^2 + \int_I x^m (u_x^2 + \frac{u^2}{x^2}) + \int_I x^m \theta_x^4 + \int_I x^m (u_x^4 + \frac{u^4}{x^4}) \\
& + \int_I x^m (w_x^4 + u_x^4 + v_x^4 + \frac{v^4}{x^4} + \frac{u^4}{x^4}) \\
\lesssim & \int_I x^m \rho\theta_t^2 + \int_I x^m \theta_x^4 + 1 \\
\lesssim & \int_I x^m \rho\theta_t^2 + \|\theta_x^2\|_{L^\infty} \int_I x^m \theta_x^2 + 1 \\
\lesssim & \int_I x^m \rho\theta_t^2 + \|\theta_x \theta_{xx}\|_{L^1} + 1 \\
\le & C \int_I x^m \rho\theta_t^2 + \frac{1}{2}\int_I x^m \theta_{xx}^2 + C.
\end{aligned}$$

This deduces

$$\int_I x^m \theta_{xx}^2 \le C \int_I x^m \rho\theta_t^2 + C. \tag{4.59}$$

Integrating (4.59) over $(0,T)$*, and using Lemma 4.19, we get*

$$\int_{Q_T} x^m \theta_{xx}^2 \le C.$$

□

Lemma 4.22 *Under the conditions of Theorem 2.1.1, we have for any* $t \in [0,T]$

$$\int_I x^m \rho_{xx}^2 + \int_{Q_T} x^m u_{xxx}^2 \le C.$$

Proof. Differentiating (4.24) with respect to x, we have

$$\begin{aligned}
\rho_{xxt} = & -\rho_{xxx}u - 3\rho_{xx}u_x - 3\rho_x u_{xx} - \rho u_{xxx} - \frac{m\rho_{xx}u}{x} - \frac{2m\rho_x u_x}{x} + \frac{2m\rho_x u}{x^2} + \frac{2m\rho u_x}{x^2} \\
& - \frac{2m\rho u}{x^3} - \frac{m\rho u_{xx}}{x}.
\end{aligned} \tag{4.60}$$

Multiplying (4.60) by $2x^m\rho_{xx}$, integrating by parts it over I, and using Hölder inequality, we have

$$\begin{aligned}\frac{d}{dt}\int_I x^m\rho_{xx}^2 =& -5\int_I x^m\rho_{xx}^2u_x - m\int_I x^{m-1}\rho_{xx}^2u - 6\int_I x^m\rho_x\rho_{xx}u_{xx} - 2\int_I x^m\rho\rho_{xx}u_{xxx}\\ & - 4m\int_I x^{m-1}\rho_x\rho_{xx}u_x + 4m\int_I x^{m-2}\rho_x\rho_{xx}u\\ & + 4m\int_I x^{m-2}\rho\rho_{xx}u_x - 4m\int_I x^{m-3}\rho\rho_{xx}u - 2m\int_I x^{m-1}\rho\rho_{xx}u_{xx}\\ =& \sum_{i=1}^{9} VII_i.\end{aligned}\tag{4.61}$$

For VII_i, i =1, 2, 3, using Cauchy inequality, Lemma 4.14 and Corollary 4.21, we have

$$\begin{aligned}VII_1+VII_2+VII_3 \lesssim& (\|u_x\|_{L^\infty}+\|u\|_{L^\infty})\int_I x^m\rho_{xx}^2 + \int_I x^m\rho_x^2u_{xx}^2 + \int_I x^m\rho_{xx}^2\\ \lesssim& \int_I x^m\rho_{xx}^2 + \|\rho_x\|_{L^\infty}^2\int_I x^m u_{xx}^2\\ \lesssim& \int_I x^m\rho_{xx}^2 + 1.\end{aligned}\tag{4.62}$$

For VII_i, i =4, 5, 6, using Cauchy inequality, Lemma 4.14 and Corollary 4.21 again, along with Lemma 4.2, we have

$$\begin{aligned}VII_4+VII_5+VII_6 \lesssim& \int_I x^m\rho_{xx}^2 + \int_I x^m u_{xxx}^2 + \int_I x^m\rho_x^2u_x^2 + \int_I x^m\rho_x^2u^2\\ \lesssim& \int_I x^m\rho_{xx}^2 + \int_I x^m u_{xxx}^2 + (\|u_x\|_{L^\infty}^2+\|u\|_{L^\infty}^2)\int_I x^m\rho_x^2\\ \lesssim& \int_I x^m\rho_{xx}^2 + \int_I x^m u_{xxx}^2 + 1.\end{aligned}\tag{4.63}$$

For VII_i, i =7, 8, 9, using Lemma 4.2 and Corollary 4.21, we have

$$\begin{aligned}VII_7+VII_8+VII_9 \lesssim& \int_I x^m\rho_{xx}^2 + \int_I x^m\rho^2u_x^2 + \int_I x^m\rho^2u^2 + \int_I x^m u_{xx}^2\rho^2\\ \lesssim& \int_I x^m\rho_{xx}^2 + 1.\end{aligned}\tag{4.64}$$

Substituting (4.62), (4.63) and (4.64) into (4.61), we have

$$\frac{d}{dt}\int_I x^m\rho_{xx}^2 \lesssim \int_I x^m\rho_{xx}^2 + \int_I x^m u_{xxx}^2 + 1.\tag{4.65}$$

Differentiating $(2.1.1)_2$ with respect to x, we have

$$\begin{aligned}\beta u_{xxx} =& \rho_xu_t + \rho u_{xt} + \rho_xuu_x + \rho u_x^2 + \rho uu_{xx} - \frac{2\rho vv_x}{x} - \frac{\rho_xv^2}{x} + \frac{\rho v^2}{x^2} + P_{xx}\\ & - \frac{m\beta(xu_{xx}-u_x)}{x^2} + \frac{m\beta(xu_x-2u)}{x^3}.\end{aligned}\tag{4.66}$$

By (4.66), we have

$$\begin{aligned}\int_I x^m u_{xxx}^2 &\lesssim \int_I x^m\rho_x^2u_t^2+\int_I x^m\rho^2u_{xt}^2+\int_I x^m\rho_x^2u^2u_x^2+\int_I x^m\rho^2u_x^4+\int_I x^m\rho^2u^2u_{xx}^2\\ &\quad+\int_I x^m\rho^2v^2v_x^2+\int_I x^m\rho_x^2v^4+\int_I x^m\rho^2v^4+\int_I x^mP_{xx}^2+\int_I x^m(u_{xx}^2+u_x^2+\frac{u^2}{x^2})\\ &\lesssim \int_I x^m\rho_x^2u_t^2+\int_I x^m\rho^2u_{xt}^2+\|u\|_{L^\infty}^2\|u_x\|_{L^\infty}^2\int_I x^m\rho_x^2+\|\rho\|_{L^\infty}^2\|u_x\|_{L^\infty}^2\int_I x^mu_x^2\\ &\quad+\|\rho\|_{L^\infty}^2\|u\|_{L^\infty}^2\int_I x^mu_{xx}^2+\|\rho\|_{L^\infty}^2\|v\|_{L^\infty}^2\int_I x^mv_x^2+\|v\|_{L^\infty}^4\int_I x^m\rho_x^2\\ &\quad+\|\rho\|_{L^\infty}\|v\|_{L^\infty}^2\int_I x^m\rho v^2+\int_I x^mP_{xx}^2+1\\ &\lesssim \int_I x^m\rho_x^2u_t^2+\int_I x^m\rho^2u_{xt}^2+\int_I x^mP_{xx}^2+1,\end{aligned} \tag{4.67}$$

where we have used Lemmas 4.2, 4.14, and Corollaries 4.16, 4.21. We need to control the right hand side of (4.67).

For the first one, we have

$$\int_I x^m\rho_x^2u_t^2 \lesssim \|u_t\|_{L^\infty}^2\int_I x^m\rho_x^2 \lesssim \int_I x^m(u_{xt}^2+\frac{u_t^2}{x^2}). \tag{4.68}$$

For the second one, we have

$$\int_I x^m\rho^2u_{xt}^2 \lesssim \|\rho\|_{L^\infty}^2\int_I x^mu_{xt}^2 \lesssim \int_I x^mu_{xt}^2. \tag{4.69}$$

For the third one, recalling $P=\rho Q+P_c(\rho)$, we have

$$\begin{aligned}\int_I x^mP_{xx}^2 &= \int_I x^m|\rho_{xx}Q+2\rho_xQ'\theta_x+\rho Q''\theta_x^2+\rho Q'\theta_{xx}+P_c''\rho_x^2+P_c'\rho_{xx}|^2\\ &\lesssim \int_I x^m\rho_{xx}^2+\int_I x^m\theta_{xx}^2+1,\end{aligned} \tag{4.70}$$

where we have used (A_3), (A_4), Lemmas 4.2, 4.14, 4.19, Corollary 4.20 and Sobolev inequality.

Substituting (4.68), (4.69) and (4.70) into (4.67), we have

$$\int_I x^mu_{xxx}^2 \lesssim \int_I x^m(u_{xt}^2+\frac{u_t^2}{x^2})+\int_I x^m\rho_{xx}^2+\int_I x^m\theta_{xx}^2+1. \tag{4.71}$$

Putting (4.71) into (4.65), we have

$$\frac{d}{dt}\int_I x^m\rho_{xx}^2 \lesssim \int_I x^m(u_{xt}^2+\frac{u_t^2}{x^2})+\int_I x^m\rho_{xx}^2+\int_I x^m\theta_{xx}^2+1. \tag{4.72}$$

By (4.72), Lemma 4.19, Corollary 4.21 and Gronwall inequality, we have

$$\int_I x^m\rho_{xx}^2 \le C. \tag{4.73}$$

Using Lemma 4.19 and Corollary 4.21 again, along with (4.71) and (4.73), we get

$$\int_{Q_T} x^mu_{xxx}^2 \le C.$$

□

Corollary 4.23 *Under the conditions of Theorem 2.1.1, we have for any* $t \in [0,T]$

$$\int_I x^m \rho_{xt}^2 + \|\rho\|_{W^{1,\infty}(Q_T)} + \|\rho_t\|_{L^\infty(Q_T)} + \int_{Q_T} x^m \rho_{tt}^2 \le C.$$

Proof. From (4.24), we have

$$\begin{aligned}\int_I x^m \rho_{xt}^2 \lesssim & \int_I x^m \rho_{xx}^2 u^2 + \int_I x^m \rho_x^2 u_x^2 + \int_I x^m \rho^2 u_{xx}^2 + \int_I x^m \rho_x^2 u^2 + \int_I x^m \rho^2 u_x^2 + \int_I x^m \rho^2 u^2 \\ \lesssim & \|u\|_{L^\infty}^2 \int_I x^m \rho_{xx}^2 + \|\rho_x\|_{L^\infty}^2 \int_I x^m u_x^2 + \|\rho\|_{L^\infty}^2 \int_I x^m u_{xx}^2 + \|u\|_{L^\infty}^2 \int_I x^m \rho_x^2 \\ & + \|\rho\|_{L^\infty}^2 \int_I x^m u_x^2 + \|\rho\|_{L^\infty} \int_I x^m \rho u^2 \le C,\end{aligned} \tag{4.74}$$

where we have used Lemmas 4.2, 4.14, Corollary 4.21, and (4.73).

By Sobolev inequality, Lemmas 4.2, 4.14, (4.73), (4.74), we have

$$\|\rho\|_{W^{1,\infty}(Q_T)} + \|\rho_t\|_{L^\infty(Q_T)} \le C. \tag{4.75}$$

Differentiating $(2.1.1)_1$ with respect to t, we have

$$\rho_{tt} = -(\rho_{xt} u + \rho_x u_t + \rho_t u_x + \rho u_{xt}) - \frac{m(\rho u_t + \rho_t u)}{x}. \tag{4.76}$$

This deduces

$$\begin{aligned}\int_I x^m \rho_{tt}^2 \lesssim & \int_I x^m \rho_{xt}^2 u^2 + \int_I x^m \rho_x^2 u_t^2 + \int_I x^m \rho_t^2 u_x^2 + \int_I x^m \rho^2 u_{xt}^2 + \int_I x^m (\rho^2 u_t^2 + \rho_t^2 u^2) \\ \lesssim & \|u\|_{L^\infty}^2 \int_I x^m \rho_{xt}^2 + \|u_t\|_{L^\infty}^2 \int_I x^m \rho_x^2 + \|u_x\|_{L^\infty}^2 \int_I x^m \rho_t^2 + \|\rho\|_{L^\infty}^2 \int_I x^m u_{xt}^2 \\ & + \|\rho\|_{L^\infty} \int_I x^m \rho u_t^2 + \|u\|_{L^\infty}^2 \int_I x^m \rho_t^2 \\ \lesssim & \int_I x^m (u_{xt}^2 + \frac{u_t^2}{x^2}) + 1,\end{aligned}$$

where we have used (4.74), Lemmas 4.2, 4.14, 4.19, and Corollary 4.21.

This combining Lemma 4.19 gives

$$\int_{Q_T} x^m \rho_{tt}^2 \le C.$$

□

Step 5: H^2 estimates of θ

Lemma 4.24 *Under the conditions of Theorem 2.1.1, we have for any* $t \in [0,T]$

$$\int_I x^m \rho \theta_t^2 + \int_{Q_T} x^m |(\kappa \theta_x)_t|^2 \le C.$$

Proof. Differentiating (4.2) with respect to t, we have

$$\rho Q' \theta_{tt} + \rho Q'' \theta_t^2 + \rho_t Q' \theta_t + (\rho u Q' \theta_x)_t + \Big(\rho \theta Q' (u_x + \frac{mu}{x})\Big)_t = (\kappa \theta_x)_{xt} + \frac{m(\kappa \theta_x)_t}{x} + \wp_t. \tag{4.77}$$

Multiplying (4.77) by $x^m \left(\int_0^\theta \kappa(\xi)\, d\xi\right)_t$ (i.e., $x^m\kappa(\theta)\theta_t$), and integrating by parts over I, we have

$$\begin{aligned}
&\frac{1}{2}\frac{d}{dt}\int_I x^m \rho Q'\kappa\theta_t^2 + \int_I x^m \left|(\kappa\theta_x)_t\right|^2 \\
=&-\frac{1}{2}\int_I x^m\rho_t Q'\kappa\theta_t^2 - \frac{1}{2}\int_I x^m\rho Q''\theta_t^3\kappa + \frac{1}{2}\int_I x^m \rho Q'\kappa'\theta_t^3 - \int_I x^m(\rho u Q'\theta_x)_t\kappa\theta_t \\
&-\int_I x^m\left(\rho\theta Q'(u_x+\frac{mu}{x})\right)_t \kappa\theta_t + \int_I x^m \wp_t\kappa\theta_t \\
=&\sum_{i=1}^{6} VIII_i.
\end{aligned}\tag{4.78}$$

For $VIII_1$, using $(2.1.1)_1$, integration by parts, (A_4), (A_5), Lemma 4.2, Corollaries 4.12, 4.20, Cauchy inequality and Poincaré inequality, we have

$$\begin{aligned}
VIII_1 =&\frac{1}{2}\int_I (x^m\rho u)_x Q'\kappa\theta_t^2 \\
=&-\frac{1}{2}\int_I x^m\rho u Q''\kappa\theta_x\theta_t^2 - \frac{1}{2}\int_I x^m\rho u Q'\kappa'\theta_x\theta_t^2 - \int_I x^m \rho u Q'\kappa\theta_t\theta_{xt} \\
\lesssim&\|\theta_x\|_{L^\infty}\int_I x^m\rho\theta_t^2 - \int_I x^m\rho u Q'\theta_t(\kappa\theta_x)_t + \int_I x^m\rho u Q'\kappa'\theta_t^2\theta_x \\
\le& C\|\theta_x\|_{L^\infty}\int_I x^m\rho\theta_t^2 + \frac{1}{10}\int_I x^m|(\kappa\theta_x)_t|^2 + C\int_I x^m\rho\theta_t^2 \\
\le& C(1+\|\theta_{xx}\|_{L^2})\int_I x^m\rho\theta_t^2 + \frac{1}{10}\int_I x^m|(\kappa\theta_x)_t|^2.
\end{aligned}\tag{4.79}$$

For $VIII_i$, i =2, 3, we have

$$\begin{aligned}
VIII_2 + VIII_3 \lesssim&\|\kappa\theta_t\|_{L^\infty}\int_I x^m\rho\theta_t^2 \\
\lesssim&(\|(\kappa\theta_t)_x\|_{L^2} + \int_I x^m\rho\kappa|\theta_t|)\int_I x^m\rho\theta_t^2 \\
\le&\frac{1}{10}\int_I x^m|(\kappa\theta_x)_t|^2 + \left(\int_I x^m\rho\theta_t^2\right)^2 + C,
\end{aligned}\tag{4.80}$$

where we have used (A_4), (A_5), Lemmas 3.1, 4.1, Corollary 4.20 and Cauchy inequality.

For $VIII_4$, we have

$$\begin{aligned}
VIII_4 =&-\int_I x^m\rho_t u Q'\theta_x\kappa\theta_t - \int_I x^m\rho u_t Q'\theta_x\kappa\theta_t - \int_I x^m\rho u Q''\theta_x\kappa\theta_t^2 - \int_I x^m\rho u Q'\theta_{xt}\kappa\theta_t \\
=&\sum_{j=1}^{4} VIII_{4,j}.
\end{aligned}\tag{4.81}$$

For $VIII_{4,1}$, using $(2.1.1)_1$, (A_4), (A_5), integration by parts, Cauchy inequality, Lemmas 4.2, 4.19, and Corollaries 4.20, 4.21, we have

$$\begin{aligned}
VIII_{4,1} =& \int_I (x^m\rho u)_x uQ'\theta_x\kappa\theta_t \\
=& -\int_I x^m\rho uu_xQ'\theta_x\kappa\theta_t - \int_I x^m\rho u^2Q''\theta_x^2\kappa\theta_t - \int_I x^m\rho u^2Q'\theta_{xx}\kappa\theta_t - \int_I x^m\rho u^2Q'\theta_x(\kappa\theta_t)_x \\
\le& C\int_I x^m\rho\theta_t^2 + C\int_I x^m\theta_x^4 + C\int_I x^m\theta_{xx}^2 + \frac{1}{20}\int_I x^m|(\kappa\theta_x)_t|^2 + C \\
\le& C\int_I x^m\rho\theta_t^2 + C\|\theta_x\|_{L^\infty}^2\int_I x^m\theta_x^2 + C\int_I x^m\theta_{xx}^2 + \frac{1}{20}\int_I x^m|(\kappa\theta_x)_t|^2 + C \\
\le& C\left(\int_I x^m\rho\theta_t^2\right)^2 + C\int_I x^m\theta_{xx}^2 + \frac{1}{20}\int_I x^m|(\kappa\theta_x)_t|^2 + C.
\end{aligned} \tag{4.82}$$

For $VIII_{4,2}$ and $VIII_{4,3}$, we have

$$\begin{aligned}
VIII_{4,2} + VIII_{4,3} \lesssim& \|\theta_x\|_{L^\infty}^2\int_I x^m\rho u_t^2 + \int_I x^m\rho\theta_t^2 + \|u\|_{L^\infty}\|\theta_x\|_{L^\infty}\int_I x^m\rho\theta_t^2 \\
\lesssim& \|\theta_x\|_{L^2}^2 + \|\theta_{xx}\|_{L^2}^2 + (1 + \|\theta_x\|_{L^2} + \|\theta_{xx}\|_{L^2})\int_I x^m\rho\theta_t^2 \\
\lesssim& \|\theta_{xx}\|_{L^2}^2 + \left(\int_I x^m\rho\theta_t^2\right)^2 + 1,
\end{aligned} \tag{4.83}$$

where we have used (A_4), (A_5), Lemma 4.19, Corollaries 4.12, 4.20, and Sobolev inequality.

For $VIII_{4,4}$, using (A_4), (A_5), Lemma 4.19, Corollaries 4.12, 4.20, and Sobolev inequality again, along with Lemma 4.2, we have

$$\begin{aligned}
VIII_{4,4} =& -\int_I x^m\rho uQ'(\kappa\theta_x)_t\theta_t + \int_I x^m\rho uQ'\kappa'\theta_x\theta_t^2 \\
\le& \frac{1}{20}\int_I x^m|(\kappa\theta_x)_t|^2 + C\left(\|\rho\|_{L^\infty}\|u\|_{L^\infty}^2 + \|u\|_{L^\infty}\|\theta_x\|_{L^\infty}\right)\int_I x^m\rho\theta_t^2 \\
\le& \frac{1}{20}\int_I x^m|(\kappa\theta_x)_t|^2 + C\int_I x^m\theta_{xx}^2 + C\left(\int_I x^m\rho\theta_t^2\right)^2 + C.
\end{aligned} \tag{4.84}$$

Putting (4.82), (4.83) and (4.84) into (4.81), we have

$$VIII_4 \le \frac{1}{10}\int_I x^m|(\kappa\theta_x)_t|^2 + C\left(\int_I x^m\rho\theta_t^2\right)^2 + C\int_I x^m\theta_{xx}^2 + C. \tag{4.85}$$

For $VIII_5$, we have

$$\begin{aligned}
VIII_5 =& -\int_I x^m\rho_t\theta Q'(u_x + \frac{mu}{x})\kappa\theta_t - \int_I x^m\rho Q'(u_x + \frac{mu}{x})\kappa\theta_t^2 \\
& -\int_I x^m\rho\theta Q''(u_x + \frac{mu}{x})\kappa\theta_t^2 - \int_I x^m\rho\theta Q'(u_{xt} + \frac{mu_t}{x})\kappa\theta_t \\
=& \sum_{j=1}^4 VIII_{5,j}.
\end{aligned} \tag{4.86}$$

For $VIII_{5,1}$, we have

$$
\begin{aligned}
VIII_{5,1} =&\int_I (x^m\rho u)_x \theta Q'(u_x+\frac{mu}{x})\kappa\theta_t \\
=&-\int_I x^m\rho u\theta_x Q'(u_x+\frac{mu}{x})\kappa\theta_t-\int_I x^m\rho u\theta\theta_x Q''(u_x+\frac{mu}{x})\kappa\theta_t \\
&-\int_I x^m\rho u\theta Q'(u_{xx}+\frac{mu_x}{x}-\frac{mu}{x^2})\kappa\theta_t-\int_I x^m\rho u\theta Q'(u_x+\frac{mu}{x})(\kappa\theta_t)_x \\
\le& C\int_I x^m\rho\theta_t^2+C\|u\|_{W^{1,\infty}}^2\int_I x^m\theta_x^2+C\int_I x^m(u_{xx}^2+u_x^2+\frac{u^2}{x^2})+\frac{1}{10}\int_I x^m|(\kappa\theta_x)_t|^2 \\
\le&\frac{1}{10}\int_I x^m|(\kappa\theta_x)_t|^2+C\int_I x^m\rho\theta_t^2+C,
\end{aligned}
\tag{4.87}
$$

where we have used $(2.1.1)_1$, integration by parts, Cauchy inequality, (A_4), (A_5), Lemmas 4.2, 4.19, and Corollaries 4.20, 4.21.

For $VIII_{5,2}$ and $VIII_{5,3}$, we have

$$
VIII_{5,2}+VIII_{5,3}\le C\int_I x^m\rho\theta_t^2. \tag{4.88}
$$

For $VIII_{5,4}$, we have

$$
VIII_{5,4}\le C\int_I x^m\rho\theta_t^2+C\int_I x^m(u_{xt}^2+\frac{u_t^2}{x^2}). \tag{4.89}
$$

Putting (4.87), (4.88) and (4.89) into (4.86), we have

$$
VIII_5\le\frac{1}{10}\int_I x^m|(\kappa\theta_x)_t|^2+C\int_I x^m\rho\theta_t^2+C\int_I x^m(u_{xt}^2+\frac{u_t^2}{x^2})+C. \tag{4.90}
$$

For $VIII_6$, recalling $\wp=\lambda(u_x+\frac{mu}{x})^2+\mu\left(w_x^2+2u_x^2+(v_x-\frac{mv}{x})^2+\frac{2mu^2}{x^2}\right)$, we have

$$
\begin{aligned}
VIII_6\lesssim&\|\kappa\theta_t\|_{L^\infty}\int_I x^m|\wp_t| \\
\lesssim&\|\kappa\theta_t\|_{L^\infty}\left((\int_I x^m u_{xt}^2)^{\frac12}+(\int_I x^m\frac{u_t^2}{x^2})^{\frac12}+(\int_I x^m w_{xt}^2)^{\frac12}+(\int_I x^m(v_{xt}^2+\frac{v_t^2}{x^2}))^{\frac12}\right) \\
\le&\frac{1}{10}\int_I x^m|(\kappa\theta_x)_t|^2+C\int_I x^m\rho\theta_t^2+C\int_I x^m u_{xt}^2+C\int_I x^m\frac{u_t^2}{x^2}+C\int_I x^m w_{xt}^2 \\
&+C\int_I x^m(v_{xt}^2+\frac{v_t^2}{x^2}),
\end{aligned}
\tag{4.91}
$$

where we have used Lemma 3.1 and Cauchy inequality.

Putting (4.79), (4.80), (4.85), (4.90) and (4.91) into (4.78), we have

$$
\begin{aligned}
&\frac{d}{dt}\int_I x^m\rho Q'\kappa\theta_t^2+\int_I x^m\left|(\kappa\theta_x)_t\right|^2 \\
\le& C\left(\int_I x^m\rho\theta_t^2\right)^2+C\int_I x^m\theta_{xx}^2+C\int_I x^m(u_{xt}^2+\frac{u_t^2}{x^2}+w_{xt}^2+v_{xt}^2+\frac{v_t^2}{x^2})+C.
\end{aligned}
\tag{4.92}
$$

By Gronwall inequality, we complete the proof of Lemma 4.24. □

Corollary 4.25 *Under the conditions of Theorem 2.1.1, we have*

$$\int_0^T \|\theta_t\|_{L^\infty}^2 \leq C.$$

Proof. By Lemma 3.1 and (A_5), we get

$$\begin{aligned}\int_0^T \|\theta_t\|_{L^\infty}^2 \lesssim & \int_0^T \|\kappa\theta_t\|_{L^\infty}^2 \\ \lesssim & \int_0^T \Big(\int_I \rho\kappa|\theta_t|\Big)^2 + \int_{Q_T} |(\kappa\theta_t)_x|^2 \\ \lesssim & \int_0^T \int_I \rho\theta_t^2 + \int_{Q_T} |(\kappa\theta_x)_t|^2 \leq C.\end{aligned}$$

□

Corollary 4.26 *Under the conditions of Theorem 2.1.1, we have*

$$\int_{Q_T} x^m\theta_{xt}^2 \leq C.$$

Proof. Since

$$\kappa\theta_{xt} = (\kappa\theta_x)_t - \kappa'\theta_t\theta_x,$$

we obtain

$$\begin{aligned}\int_{Q_T} x^m\theta_{xt}^2 \lesssim & \int_{Q_T} x^m\kappa^2\theta_{xt}^2 \\ \lesssim & \int_{Q_T} x^m\,|(\kappa\theta_x)_t|^2 + \int_{Q_T} x^m(\kappa')^2\theta_t^2\theta_x^2 \\ \lesssim & 1 + \int_0^T \sup_{x\in I}\theta_t^2 \int_I x^m\theta_x^2 \leq C,\end{aligned}$$

where we have used (A_5), Lemmas 4.19, 4.24, and Corollaries 4.20, 4.25.

□

Corollary 4.27 *Under the conditions of Theorem 2.1.1, we have for any* $t \in [0,T]$

$$\|\theta\|_{W^{1,\infty}(Q_T)} + \int_I x^m\theta_{xx}^2 + \int_{Q_T} x^m\theta_{xxx}^2 \leq C.$$

Proof. By (4.59) and Lemma 4.24, we get

$$\int_I x^m\theta_{xx}^2 \leq C.$$

This, along with Lemma 4.19, Corollary 4.20 and Sobolev inequality, gives

$$\|\theta\|_{W^{1,\infty}(Q_T)} \leq C.$$

Differentiating (4.2) w.r.t. x, we have

$$\begin{aligned}\kappa\theta_{xxx} = & -(\frac{m\kappa\theta_x}{x})_x - 3\kappa'\theta_x\theta_{xx} - \kappa''\theta_x^3 - \wp_x + \rho Q'\theta_{xt} + \rho_x Q'\theta_t + \rho Q''\theta_x\theta_t \\ & + (\rho u Q'\theta_x)_x + \Big(\rho\theta Q'(u_x + \frac{mu}{x})\Big)_x.\end{aligned} \tag{4.93}$$

This deduces

$$\begin{aligned}\int_I x^m\theta_{xxx}^2 &\lesssim \int_I x^m\rho^2\theta_{xt}^2+\int_I x^m\rho_x^2\theta_t^2+1\\ &\lesssim \int_I x^m\theta_{xt}^2+\sup_{x\in I}\theta_t^2+1.\end{aligned}\tag{4.94}$$

Integrating (4.94) over $(0,T)$, and using Corollary 4.25 and Corollary 4.26, we get

$$\int_{Q_T} x^m\theta_{xxx}^2\le C.$$

□

Constructing a sequence of approximate initial data and passing to the limits: Denote $\rho_0^\varepsilon=\rho_0+\varepsilon$ and $\theta_0^\varepsilon=\theta_0+\varepsilon$. u_0^ε is the solution to the following elliptic problem:

$$\begin{cases}\beta(u_{0xx}^\varepsilon+\frac{mu_{0x}^\varepsilon}{x}-\frac{mu_0^\varepsilon}{x^2})-P_x(\rho_0^\varepsilon,\theta_0^\varepsilon)=\sqrt{\rho}_0g_1,\ x\in I,\\ u_0^\varepsilon|_{x=a,b}=0.\end{cases}\tag{4.95}$$

Since $\rho_0^\varepsilon=\rho_0+\varepsilon\in H^2$, $\theta_0^\varepsilon=\theta_0+\varepsilon\in H^2$ and $g_1\in L^2$, we obtain by using the standard elliptic estimates

$$\begin{cases}u_0^\varepsilon\in H^2\cap H_0^1,\\ \|u_0^\varepsilon-u_0\|_{H^2}\le C\varepsilon.\end{cases}\tag{4.96}$$

Consider (2.1.1)-(2.1.3) with initial data replaced by (ρ_0^ε, u_0^ε, v_0, w_0, θ_0^ε), we get a global solution (ρ^ε, u^ε, v^ε, w^ε, θ^ε) to (2.1.1)-(2.1.3) for each $\varepsilon>0$. These *a priori* estimates in the section are valid for (ρ^ε, u^ε, v^ε, w^ε, θ^ε). Then taking the limits $\varepsilon\to 0^+$ (taking subsequence if necessary), and using Lemma 3.5, we get a solution denoted by (ρ, u, v, w, θ) to (2.1.1)-(2.1.3) with regularities as in Theorem 2.1.1. The uniqueness of the solutions can be done by the standard methods, see for instance [4] and references therein. We end the proof of Theorem 2.1.1. □

Remark 4.28 *The global existence of (ρ^ε, u^ε, v^ε, w^ε, θ^ε) can be done by local existence (using the similar arguments as in [4])+ some a priori estimates globally in time throughout the section.*

5 Proof of Theorem 2.1.6

In the section, we denote by C a generic constant depending only on $\|(\rho_0,u_0,v_0,w_0,\theta_0)\|_{H^3}$, $\|g_i\|_{L^2}$, $\|(\sqrt{\rho_0}g_i)_x\|_{L^2}$ $(i=1,2,3,4)$, T, λ, μ, a, b, and some other known constants, but independent of the solutions and the lower bounds of the density.

Under the conditions of Theorem 2.1.6, all those *a priori* estimates in section 3 are also satisfied in this section. Following the quite similar strategies with the proof of Theorem 2.1.1, to prove Theorem 2.1.6, it suffices to get H^3-*a-priori*-estimates of the classical solutions (ρ, u, v, w, θ) as in Theorem 2.1.6 with $\inf\limits_{(x,t)\in Q_T}\rho>0$.

To begin with, we get a $W^{1,\infty}(Q_T)$-estimate of $\sqrt{\rho}$ which was obtained in [8, 7] as a crucial estimate to get H^4-estimate of u.

Lemma 5.1 *Under the conditions of Theorem 2.1.6, we have*

$$\|(\sqrt{\rho})_x\|_{L^\infty(Q_T)}+\|(\sqrt{\rho})_t\|_{L^\infty(Q_T)}\le C.$$

Proof. Multiplying $(2.1.1)_1$ by $\frac{1}{2\sqrt{\rho}}$, we have

$$(\sqrt{\rho})_t+\frac{mx^{-1}}{2}\sqrt{\rho}u+\frac{1}{2}\sqrt{\rho}u_x+(\sqrt{\rho})_x u=0. \tag{5.1}$$

Differentiating (5.1) with respect to x, we get

$$(\sqrt{\rho})_{xt}+\frac{mx^{-1}}{2}(\sqrt{\rho})_x u+\frac{mx^{-1}}{2}\sqrt{\rho}u_x-\frac{m\sqrt{\rho}u}{2x^2}+\frac{3}{2}(\sqrt{\rho})_x u_x+\frac{1}{2}\sqrt{\rho}u_{xx}+(\sqrt{\rho})_{xx}u=0.$$

Denote $h=(\sqrt{\rho})_x$, we have

$$h_t+h_x u+h(\frac{mx^{-1}u}{2}+\frac{3u_x}{2})+\frac{mx^{-1}}{2}\sqrt{\rho}u_x-\frac{m\sqrt{\rho}u}{2x^2}+\frac{1}{2}\sqrt{\rho}u_{xx}=0,$$

which implies

$$\begin{aligned}&\frac{d}{dt}\left\{h\exp\left[\int_0^t(\frac{mx^{-1}u}{2}+\frac{3u_x}{2})(x(\tau,y),\tau)d\tau\right]\right\}\\ =&-\left(\frac{mx^{-1}\sqrt{\rho}u_x}{2}-\frac{m\sqrt{\rho}u}{2x^2}+\frac{\sqrt{\rho}u_{xx}}{2}\right)\\ &\times\exp\left[\int_0^t\left(\frac{mx^{-1}u}{2}+\frac{3u_x}{2}\right)(x(\tau,y),\tau)d\tau\right],\end{aligned} \tag{5.2}$$

where $x(t,y)$ satisfies

$$\begin{cases}\frac{dx(t,y)}{dt}=u(x(t,y),t),\ 0\le t<s,\\ x(s,y)=y.\end{cases}$$

Integrating (5.2) over $(0,s)$, we get

$$\begin{aligned}h(y,s)=&\exp\left(-\int_0^s\left(\frac{mx^{-1}u}{2}+\frac{3u_x}{2}\right)(x(\tau,y),\tau)d\tau\right)h(x(0,y),0)\\ &-\int_0^s\Big[(\frac{mx^{-1}\sqrt{\rho}u_x}{2}-\frac{m\sqrt{\rho}u}{2x^2}+\frac{\sqrt{\rho}u_{xx}}{2})\\ &\qquad\times\exp\left(\int_s^t\left(\frac{mx^{-1}u}{2}+\frac{3u_x}{2}\right)(x(\tau,y),\tau)d\tau\right)\Big]dt.\end{aligned}$$

This implies

$$\|(\sqrt{\rho})_x\|_{L^\infty(Q_T)}\le C. \tag{5.3}$$

From (5.1) and (5.3), we get

$$\|(\sqrt{\rho})_t\|_{L^\infty(Q_T)}\le C.$$

The proof of Lemma 5.1 is complete. □

Lemma 5.2 *Under the conditions of Theorem 2.1.6, we have for any* $t\in[0,T]$

$$\int_I x^m\rho^2\left|(\kappa\theta_x)_t\right|^2+\int_{Q_T}x^m\rho^3\theta_{tt}^2\le C.$$

Proof. Multiplying (4.77) by $x^m\rho^\alpha(\kappa\theta_t)_t$ (i.e. $x^m\rho^\alpha\kappa\theta_{tt}+x^m\rho^\alpha\kappa'\theta_t^2$, where $\alpha>0$ is to be decided later), and integrating by parts over I, we have

$$\begin{aligned}&\int_I x^m\rho^{\alpha+1}\kappa Q'\theta_{tt}^2+\frac{1}{2}\frac{d}{dt}\int_I x^m\rho^\alpha\left|(\kappa\theta_x)_t\right|^2\\ =&\frac{\alpha}{2}\int_I x^m\rho^{\alpha-1}\rho_t\left|(\kappa\theta_x)_t\right|^2-\alpha\int_I x^m\rho^{\alpha-1}\rho_x\kappa\theta_{tt}(\kappa\theta_x)_t-\alpha\int_I x^m\rho^{\alpha-1}\rho_x\kappa'\theta_t^2(\kappa\theta_x)_t\\ &+\int_I x^m\rho^\alpha\wp_t(\kappa\theta_{tt}+\kappa'\theta_t^2)-\int_I x^m\rho^{\alpha+1}Q'\kappa'\theta_t^2\theta_{tt}-\int_I x^m\rho^\alpha\left[\rho Q''\theta_t^2+\rho_tQ'\theta_t\right]\left(\kappa\theta_{tt}+\kappa'\theta_t^2\right)\\ &-\int_I x^m\rho^\alpha(\rho uQ'\theta_x)_t\left(\kappa\theta_{tt}+\kappa'\theta_t^2\right)-\int_I x^m\rho^\alpha\left(\rho\theta Q'(u_x+\frac{mu}{x})\right)_t\left(\kappa\theta_{tt}+\kappa'\theta_t^2\right)=\sum_{i=1}^{8}VIV_i.\end{aligned}\tag{5.4}$$

For VIV_2, using (A_4), (A_5), Corollary 4.20 and Cauchy inequality, we have

$$\begin{aligned}VIV_2\leq&\frac{1}{10}\int_I x^m\rho^{\alpha+1}\kappa Q'\theta_{tt}^2+C\int_I x^m\rho^{\alpha-3}\rho_x^2|(\kappa\theta_x)_t|^2\\ \leq&\frac{1}{10}\int_I x^m\rho^{\alpha+1}\kappa Q'\theta_{tt}^2+C\|\rho^{\alpha-3}\rho_x^2\|_{L^\infty}\int_I x^m|(\kappa\theta_x)_t|^2.\end{aligned}\tag{5.5}$$

For VIV_1 and VIV_3, we have

$$\begin{aligned}VIV_1+VIV_3\leq&C\|\rho^{\alpha-1}\rho_t\|_{L^\infty}\int_I x^m\left|(\kappa\theta_x)_t\right|^2+C\|\theta_t\|_{L^\infty}^2\int_I x^m\rho^\alpha|(\kappa\theta_x)_t|^2\\ &+C\|\theta_t\|_{L^\infty}^2\int_I x^m\rho^{\alpha-2}\rho_x^2.\end{aligned}\tag{5.6}$$

For VIV_4, we have

$$\begin{aligned}VIV_4\leq&\frac{1}{10}\int_I x^m\rho^{\alpha+1}\kappa Q'\theta_{tt}^2+C\|\rho^{\alpha-1}\|_{L^\infty}\int_I x^m\wp_t^2+C\int_I x^m\rho^{\alpha+1}\theta_t^4\\ \leq&\frac{1}{10}\int_I x^m\rho^{\alpha+1}\kappa Q'\theta_{tt}^2+C\|\rho^{\alpha-1}\|_{L^\infty}\int_I x^m(u_{xt}^2+\frac{u_t^2}{x^2}+w_{xt}^2+v_{xt}^2+\frac{v_t^2}{x^2})+C\|\theta_t\|_{L^\infty}^2,\end{aligned}\tag{5.7}$$

where we have used Cauchy inequality, Corollaries 4.16, 4.18, 4.21 and Lemmas 4.2, 4.24.

For VIV_5 and VIV_6, using (A_4), (A_5), Cauchy inequality, Corollary 4.20, and Lemmas 4.2, 4.24, we have

$$\begin{aligned}VIV_5+VIV_6=&-\int_I x^m\rho^{\alpha+1}Q'\kappa'\theta_t^2\theta_{tt}-\int_I x^m\rho^\alpha\left[\rho Q''\theta_t^2+\rho_tQ'\theta_t\right]\kappa\theta_{tt}\\ &-\int_I x^m\rho^\alpha\left[\rho Q''\theta_t^2+\rho_tQ'\theta_t\right]\kappa'\theta_t^2\\ \leq&\frac{1}{10}\int_I x^m\rho^{\alpha+1}\kappa Q'\theta_{tt}^2+C\int_I x^m\rho^{\alpha+1}\theta_t^4+C\int_I x^m\rho^{\alpha-1}\rho_t^2\theta_t^2\\ \leq&\frac{1}{10}\int_I x^m\rho^{\alpha+1}\kappa Q'\theta_{tt}^2+C\|\theta_t\|_{L^\infty}^2(1+\int_I x^m\rho^{\alpha-1}\rho_t^2)+C.\end{aligned}\tag{5.8}$$

For VIV_7, using (A_4), (A_5), Cauchy inequality, and Lemmas 4.2, 4.24 again, along with

Corollaries 4.12, 4.27 and Lemma 4.19, we have

$$\begin{aligned}
VIV_7 =& -\int_I x^m\rho^\alpha\rho_t uQ'\theta_x\left(\kappa\theta_{tt}+\kappa'\theta_t^2\right)-\int_I x^m\rho^{\alpha+1}u_tQ'\theta_x\left(\kappa\theta_{tt}+\kappa'\theta_t^2\right)\\
&-\int_I x^m\rho^{\alpha+1}uQ''\theta_t\theta_x\left(\kappa\theta_{tt}+\kappa'\theta_t^2\right)-\int_I x^m\rho^{\alpha+1}uQ'\theta_{xt}\left(\kappa\theta_{tt}+\kappa'\theta_t^2\right)\\
\le&\frac{1}{10}\int_I x^m\rho^{\alpha+1}\kappa Q'\theta_{tt}^2+C\int_I x^m\rho^{\alpha-1}\rho_t^2+C\int_I x^m\rho^{\alpha+1}\theta_t^4+C\int_I x^m\rho^{\alpha+1}u_t^2\\
&+C\int_I x^m\rho^{\alpha+1}\theta_t^2+C\int_I x^m\rho^{\alpha+1}\theta_{xt}^2\\
\le&\frac{1}{10}\int_I x^m\rho^{\alpha+1}\kappa Q'\theta_{tt}^2+C\int_I x^m\rho^{\alpha-1}\rho_t^2+C\|\theta_t\|_{L^\infty}^2+C\int_I x^m\theta_{xt}^2+C.
\end{aligned}\tag{5.9}$$

Similarly, for VIV_8, we have

$$\begin{aligned}
VIV_8 =& -\int_I x^m\rho^\alpha\rho_t\theta Q'(u_x+\frac{mu}{x})\left(\kappa\theta_{tt}+\kappa'\theta_t^2\right)-\int_I x^m\rho^{\alpha+1}\theta_tQ'(u_x+\frac{mu}{x})\left(\kappa\theta_{tt}+\kappa'\theta_t^2\right)\\
&-\int_I x^m\rho^{\alpha+1}\theta Q''\theta_t(u_x+\frac{mu}{x})\left(\kappa\theta_{tt}+\kappa'\theta_t^2\right)-\int_I x^m\rho^{\alpha+1}\theta Q'(u_{xt}+\frac{mu_t}{x})\left(\kappa\theta_{tt}+\kappa'\theta_t^2\right)\\
\le&\frac{1}{10}\int_I x^m\rho^{\alpha+1}\kappa Q'\theta_{tt}^2+C\int_I x^m\rho^{\alpha-1}\rho_t^2+C\int_I x^m\rho^{\alpha+1}\theta_t^4\\
&+C\int_I x^m\rho^{\alpha+1}\theta_t^2+C\int_I x^m\rho^{\alpha+1}(u_{xt}^2+\frac{u_t^2}{x^2})\\
\le&\frac{1}{10}\int_I x^m\rho^{\alpha+1}\kappa Q'\theta_{tt}^2+C\int_I x^m\rho^{\alpha-1}\rho_t^2+C\|\theta_t\|_{L^\infty}^2+C\int_I x^m(u_{xt}^2+\frac{u_t^2}{x^2})+C.
\end{aligned}\tag{5.10}$$

Putting (5.5), (5.6), (5.7), (5.8), (5.9) and (5.10) into (5.4), we have

$$\begin{aligned}
&\int_I x^m\rho^{\alpha+1}\kappa Q'\theta_{tt}^2+\frac{d}{dt}\int_I x^m\rho^\alpha\left|(\kappa\theta_x)_t\right|^2\\
\le& C\|\theta_t\|_{L^\infty}^2\int_I x^m\rho^\alpha|(\kappa\theta_x)_t|^2+C(\|\rho^{\alpha-3}\rho_x^2\|_{L^\infty}+\|\rho^{\alpha-1}\rho_t\|_{L^\infty})\int_I x^m\left|(\kappa\theta_x)_t\right|^2\\
&+C(1+\int_I x^m\rho^{\alpha-1}\rho_t^2+\int_I x^m\rho^{\alpha-2}\rho_x^2)\|\theta_t\|_{L^\infty}^2+C\int_I x^m\rho^{\alpha-1}\rho_t^2+C\int_I x^m\theta_{xt}^2\\
&+C\|\rho^{\alpha-1}\|_{L^\infty}\int_I x^m(u_{xt}^2+\frac{u_t^2}{x^2}+w_{xt}^2+v_{xt}^2+\frac{v_t^2}{x^2})+C\int_I x^m(u_{xt}^2+\frac{u_t^2}{x^2})+C.
\end{aligned}\tag{5.11}$$

Claim:

$$\int_I x^m\rho^\alpha\left|(\kappa\theta_x)_t\right|^2+\int_{Q_T}x^m\rho^{\alpha+1}\kappa Q'\theta_{tt}^2\le C,\tag{5.12}$$

if

$$\|\rho^{\alpha-3}\rho_x^2\|_{L^\infty}+\|\rho^{\alpha-1}\rho_t\|_{L^\infty}+\|\rho^{\alpha-1}\|_{L^\infty}\le C.\tag{5.13}$$

In fact, if (5.13) is satisfied, (5.12) will be obtained by using (5.11), Corollaries 4.25, 4.26, Lemmas 4.15, 4.17, 4.19, 4.24, (2.1.4) and Gronwall inequality.

To ensure that (5.13) is valid, the restriction $\alpha\ge 3$ seems necessary. While, to get $L_t^\infty H_x^3$-estimate of θ (Corollary 5.3), by (4.2), α in (5.12) should satisfy $\alpha\le 2$. Assume $\alpha\in[1,2]$, to get (5.13), it suffices to prove

$$\|\rho^{\alpha-3}\rho_x^2\|_{L^\infty}\le C.$$

In fact,

$$\|\rho^{\alpha-3}\rho_x^2\|_{L^\infty} = 4\left\|\rho^{\alpha-2}|(\sqrt{\rho})_x|^2\right\|_{L^\infty}.$$

Assume $\alpha = 2$, and use Lemma 5.1, (5.13) is to be obtained. This means that (5.12) is obtained for $\alpha = 2$. The proof of Lemma 5.2 is complete. □

Corollary 5.3 *Under the conditions of Theorem 2.1.6, we have for any* $t \in [0,T]$

$$\int_I x^m \left(\theta_{xxx}^2 + \rho^2\theta_{xt}^2\right) \le C.$$

Proof. A direct calculation gives

$$\rho\kappa\theta_{xt} = \rho(\kappa\theta_x)_t - \rho\kappa'\theta_t\theta_x,$$

which implies

$$\begin{aligned}\int_I x^m\rho^2\theta_{xt}^2 \le& C\int_I x^m\rho^2\left|(\kappa\theta_x)_t\right|^2 + C\|\theta_x\|_{L^\infty}^2\int_I x^m\rho\theta_t^2\\ \le& C,\end{aligned}\tag{5.14}$$

where we have used (A_5), Lemma 4.2, Lemma 4.24, Lemma 5.2 and Corollary 4.27.

From the first inequality of (4.94), using Lemma 4.24, (5.14) and Lemma 5.1, we obtain

$$\begin{aligned}\int_I x^m\theta_{xxx}^2 \le& C\int_I x^m\rho^2\theta_{xt}^2 + C\int_I x^m\rho_x^2\theta_t^2 + C\\ \le& C\int_I x^m\rho^2\theta_{xt}^2 + C\int_I x^m\rho|(\sqrt{\rho})_x|^2\theta_t^2 + C \le C.\end{aligned}$$

□

Lemma 5.4 *Under the conditions of Theorem 2.1.6, we have for any* $t \in [0,T]$

$$\int_I x^m\rho^2u_{xt}^2 + \int_{Q_T} x^m\rho^3u_{tt}^2 \le C.$$

Proof. Similar to Lemma 5.2, multiplying (4.42) by $x^m\rho^2u_{tt}$, and integrating by parts over I, we have

$$\begin{aligned}&\int_I x^m\rho^3u_{tt}^2 + \frac{\beta}{2}\frac{d}{dt}\int_I x^m\rho^2u_{xt}^2\\ =&\beta\int_I x^m\rho\rho_tu_{xt}^2 - 2\beta\int_I x^m\rho\rho_xu_{xt}u_{tt} - \int_I x^m\rho^2u_{tt}(\rho_tu_t + \rho_tuu_x + \rho u_tu_x + \rho uu_{xt} - \frac{\rho_tv^2}{x} - \frac{2\rho vv_t}{x})\\ &-\beta\int_I x^m\rho^2u_{tt}\frac{mu_t}{x^2} - \int_I x^m\rho^2u_{tt}P_{xt}\\ =&\sum_{i=1}^5 VV_i.\end{aligned}\tag{5.15}$$

For VV_1 and VV_2, using Corollary 4.23, Lemma 5.1 and Cauchy inequality, we have

$$\begin{aligned}VV_1 + VV_2 \le& \|\rho\|_{L^\infty}\|\rho_t\|_{L^\infty}\int_I x^mu_{xt}^2 - 4\beta\int_I x^m\rho^{\frac{3}{2}}(\sqrt{\rho})_xu_{xt}u_{tt}\\ \le& C\int_I x^mu_{xt}^2 + \frac{1}{8}\int_I x^m\rho^3u_{tt}^2.\end{aligned}\tag{5.16}$$

For VV_3, we have

$$\begin{aligned}
VV_3 \leq & \frac{1}{8}\int_I x^m\rho^3 u_{tt}^2 + C\int_I x^m \rho\rho_t^2 u_t^2 + C\int_I x^m\rho\rho_t^2 u^2 u_x^2 + C\int_I x^m \rho^3 u_t^2 u_x^2 \\
& + C\int_I x^m \rho^3 u^2 u_{xt}^2 + C\int_I x^m \frac{\rho\rho_t^2 v^4}{x^2} + C\int_I x^m \frac{\rho^3 v^2 v_t^2}{x^2} \\
\leq & \frac{1}{8}\int_I x^m\rho^3 u_{tt}^2 + C\|\rho_t\|_{L^\infty}^2 \int_I x^m \rho u_t^2 + C\|\rho\|_{L^\infty}\|\rho_t\|_{L^\infty}^2\|u\|_{L^\infty}^2 \int_I x^m u_x^2 \\
& + C\|u_x\|_{L^\infty}^2\|\rho\|_{L^\infty}^2 \int_I x^m\rho u_t^2 + C\|\rho\|_{L^\infty}^3\|u\|_{L^\infty}^2 \int_I x^m u_{xt}^2 \\
& + C\|\rho_t\|_{L^\infty}^2\|v\|_{L^\infty}^2 \int_I x^m \rho v^2 + C\|v\|_{L^\infty}^2\|\rho\|_{L^\infty}^2 \int_I x^m\rho v_t^2 \\
\leq & \frac{1}{8}\int_I x^m\rho^3 u_{tt}^2 + C\int_I x^m u_{xt}^2 + C,
\end{aligned} \tag{5.17}$$

where we have used Cauchy inequality, Corollaries 4.8, 4.21, 4.23, and Lemmas 4.15, 4.19.

For VV_4, using Cauchy inequality and Lemma 4.19, we have

$$VV_4 \leq \frac{1}{8}\int_I x^m \rho^3 u_{tt}^2 + C\int_I x^m\rho u_t^2 \leq \frac{1}{8}\int_I x^m\rho^3 u_{tt}^2 + C. \tag{5.18}$$

For VV_5, recalling $P = \rho Q + P_c$, we have

$$\begin{aligned}
VV_5 = & -\int_I x^m \rho^2 u_{tt}\left(\rho_{xt}Q + \rho_x Q'\theta_t + \rho_t Q'\theta_x + \rho Q''\theta_t\theta_x + \rho Q'\theta_{xt} + P_c'\rho_{xt} + P_c''\rho_x\rho_t\right) \\
\leq & \frac{1}{8}\int_I x^m\rho^3 u_{tt}^2 + C\int_I x^m\rho\rho_{xt}^2 Q^2 + \int_I x^m \rho\rho_x^2 (Q')^2\theta_t^2 + \int_I x^m\rho\rho_t^2(Q')^2\theta_x^2 \\
& + \int_I x^m\rho^3 (Q'')^2\theta_t^2\theta_x^2 + \int_I x^m \rho^3 (Q')^2\theta_{xt}^2 + \int_I x^m\rho(P_c')^2\rho_{xt}^2 + \int_I x^m \rho (P_c'')^2\rho_x^2\rho_t^2 \\
\leq & \frac{1}{8}\int_I x^m\rho^3 u_{tt}^2 + C,
\end{aligned} \tag{5.19}$$

where we have used Cauchy inequality, (A_3), (A_4), Corollaries 4.23, 4.27, 5.3, and Lemma 4.24.

Putting (5.16), (5.17), (5.18) and (5.19) into (5.15), we have

$$\int_I x^m\rho^3 u_{tt}^2 + \beta\frac{d}{dt}\int_I x^m\rho^2 u_{xt}^2 \leq C\int_I x^m u_{xt}^2 + C. \tag{5.20}$$

By (5.20), Lemma 4.19 and Gronwall inequality, we complete the proof of Lemma 5.4. □

Corollary 5.5 *Under the conditions of Theorem 2.1.6, we have for any* $t\in[0,T]$

$$\int_I x^m u_{xxx}^2 \leq C.$$

Proof. By (4.67), (4.70), Lemmas 4.19, 4.22, 5.1, 5.4, Corollaries 4.27, we get

$$\begin{aligned}
\int_I x^m u_{xxx}^2 \leq & C\int_I x^m\rho_x^2 u_t^2 + C\int_I x^m\rho^2 u_{xt}^2 + C\int_I x^m P_{xx}^2 + C \\
\leq & C\int_I x^m \rho|(\sqrt{\rho})_x|^2 u_t^2 + C \leq C.
\end{aligned} \tag{5.21}$$

□

Lemma 5.6 *Under the conditions of Theorem 2.1.6, we have for any* $t \in [0,T]$

$$\int_I x^m\rho^2(v_{xt}^2+w_{xt}^2)+\int_{Q_T} x^m\rho^3(v_{tt}^2+w_{tt}^2)\le C.$$

Proof. Multiplying (4.36) by $x^m\rho^2 v_{tt}$, integrating by parts over I, we have

$$\begin{aligned}
&\int_I x^m\rho^3 v_{tt}^2+\frac{\mu}{2}\frac{d}{dt}\int_I x^m\rho^2 v_{xt}^2\\
=&\mu\int_I x^m\rho\rho_t v_{xt}^2-2\mu\int_I x^m\rho\rho_x v_{xt}v_{tt}-\int_I x^m\rho^2 v_{tt}(\rho_t v_t+\rho_t u v_x+\rho u_t v_x+\rho u v_{xt})\\
&-\int_I x^m\rho^2 v_{tt}(x^{-1}\rho_t uv+x^{-1}\rho u_t v+x^{-1}\rho u v_t)-\mu\int_I x^m\rho^2 v_{tt}\frac{mv_t}{x^2}\\
=&\sum_{i=1}^5 VVI_i.
\end{aligned}\tag{5.22}$$

For VVI_1 and VVI_2, using Cauchy inequality, Corollary 4.23 and Lemma 5.1, we have

$$\begin{aligned}
VVI_1+VVI_2\le&\|\rho\|_{L^\infty}\|\rho_t\|_{L^\infty}\int_I x^m v_{xt}^2-4\mu\int_I x^m\rho^{\frac{3}{2}}(\sqrt{\rho})_x v_{xt}v_{tt}\\
\le&\frac{1}{8}\int_I x^m\rho^3 v_{tt}^2+C\int_I x^m v_{xt}^2.
\end{aligned}\tag{5.23}$$

For VVI_3, using Cauchy inequality, Corollary 4.23 again, together with Corollaries 4.12, 4.16, and Lemmas 4.15, 4.19, we have

$$\begin{aligned}
VVI_3\le&\frac{1}{8}\int_I x^m\rho^3 v_{tt}^2+C\|\rho_t\|_{L^\infty}^2\int_I x^m\rho v_t^2+C\|\rho\|_{L^\infty}\|\rho_t\|_{L^\infty}^2\|u\|_{L^\infty}^2\int_I x^m v_x^2\\
&+C\|v_x\|_{L^\infty}^2\|\rho\|_{L^\infty}^2\int_I x^m\rho u_t^2+C\|\rho\|_{L^\infty}^3\|u\|_{L^\infty}^2\int_I x^m v_{xt}^2\\
\le&\frac{1}{8}\int_I x^m\rho^3 v_{tt}^2+C\int_I x^m v_{xt}^2+C.
\end{aligned}\tag{5.24}$$

Similarly, for VVI_4 and VVI_5, we have

$$\begin{aligned}
VVI_4\le&\frac{1}{8}\int_I x^m\rho^3 v_{tt}^2+C\|\rho\|_{L^\infty}\|u\|_{L^\infty}^2\|v\|_{L^\infty}^2\int_I x^m\rho_t^2+C\|\rho\|_{L^\infty}^2\|v\|_{L^\infty}^2\int_I x^m\rho u_t^2\\
&+C\|\rho\|_{L^\infty}^2\|u\|_{L^\infty}^2\int_I x^m\rho v_t^2\\
\le&\frac{1}{8}\int_I x^m\rho^3 v_{tt}^2+C;
\end{aligned}\tag{5.25}$$

$$VVI_5\le\frac{1}{8}\int_I x^m\rho^3 v_{tt}^2+C\int_I x^m\rho v_t^2\le\frac{1}{8}\int_I x^m\rho^3 v_{tt}^2+C.\tag{5.26}$$

Putting (5.23), (5.24), (5.25) and (5.26) into (5.22), and using Lemma 4.15, Gronwall inequality, we have

$$\int_I x^m\rho^2 v_{xt}^2+\int_{Q_T} x^m\rho^3 v_{tt}^2\le C.\tag{5.27}$$

Similarly, we get

$$\int_I x^m\rho^2 w_{xt}^2+\int_{Q_T} x^m\rho^3 w_{tt}^2\le C.$$

□

Corollary 5.7 *Under the conditions of Theorem 2.1.6, we have for any* $t \in [0,T]$

$$\int_I x^m (v_{xxx}^2 + w_{xxx}^2) + \int_{Q_T} (v_{xxxx}^2 + w_{xxxx}^2) \le C.$$

Proof. Similar to Corollary 5.5, we get

$$\int_I x^m (v_{xxx}^2 + w_{xxx}^2) \le C.$$

By (4.36), Corollaries 4.12, 4.16, 4.23, and Lemmas 4.15, 4.19, 5.6, we have

$$\int_{Q_T} \rho v_{xxt}^2 \le C.$$

This together with $(2.1.1)_3$ gives

$$\int_{Q_T} v_{xxxx}^2 \le C.$$

Similarly, we can get

$$\int_{Q_T} w_{xxxx}^2 \le C.$$

□

Lemma 5.8 *Under the conditions of Theorem 2.1.6, we have for any* $t \in [0,T]$

$$\int_I x^m \rho_{xxx}^2 + \int_{Q_T} x^m (u_{xxxx}^2 + \theta_{xxxx}^2) \le C.$$

Proof. Differentiating (4.60) with respect to x, multiplying it by $x^m \rho_{xxx}$, and integrating by parts over I, we have

$$\begin{aligned}\frac{1}{2}\frac{d}{dt}\int_I x^m \rho_{xxx}^2 = &\int_I x^m \rho_{xxx}\Big[- m x^{-1} \rho u_{xxx} - 3m x^{-1} \rho_x u_{xx} - 3m x^{-1} \rho_{xx} u_x + \frac{3m\rho u_{xx}}{x^2} \\ &- \frac{6m\rho u_x}{x^3} - m x^{-1} \rho_{xxx} u + \frac{3m\rho_{xx} u}{x^2} + \frac{6m\rho_x u_x}{x^2} - \frac{6m\rho_x u}{x^3} + \frac{6m\rho u}{x^4} - 4\rho_{xxx} u_x - 6\rho_{xx} u_{xx} \\ &- 4\rho_x u_{xxx}\Big] + \frac{1}{2}\int_I x^m \rho_{xxx}^2 u_x + \frac{1}{2}\int_I m x^{m-1} \rho_{xxx}^2 u - \int_I x^m \rho \rho_{xxx} u_{xxxx}.\end{aligned}$$

By Sobolev inequality, Cauchy inequality, Corollaries 4.21, 4.23, 5.5, and Lemma 4.22, we get

$$\frac{d}{dt}\int_I x^m \rho_{xxx}^2 \le C\int_I x^m \rho_{xxx}^2 + C\int_I x^m u_{xxxx}^2 + C. \tag{5.28}$$

Differentiating (4.66) with respect to x, we have

$$\begin{aligned}\beta u_{xxxx} =&\rho_{xx} u_t + 2\rho_x u_{xt} + \rho u_{xxt} + (\rho_x u u_x + \rho u_x^2 + \rho u u_{xx})_x - (\frac{2\rho v v_x}{x} + \frac{\rho_x v^2}{x} - \frac{\rho v^2}{x^2})_x \\ &+ P_{xxx} - \frac{m\beta u_{xxx}}{x} + \frac{3m\beta u_{xx}}{x^2} - \frac{6m\beta u_x}{x^3} + \frac{6m\beta u}{x^4}.\end{aligned} \tag{5.29}$$

Using Lemma 4.22, Corollaries 4.16, 4.21, 4.23, 4.27, 5.3, 5.5, and (A_3), (A_4), we obtain

$$\begin{aligned}\int_I x^m u_{xxxx}^2 \le& C\int_I x^m |P_{xxx}|^2 + C\int_I x^m (\rho u_{xxt}^2 + u_{xt}^2) + C \\ \le& C\int_I x^m \rho_{xxx}^2 + C\int_I x^m (\rho u_{xxt}^2 + u_{xt}^2) + C.\end{aligned} \tag{5.30}$$

Substituting (5.30) into (5.28), and using (4.42), Lemmas 4.19, 5.4, Gronwall inequality, we obtain

$$\int_I x^m \rho_{xxx}^2 \le C. \tag{5.31}$$

By (5.30) and (5.31), we have

$$\int_{Q_T} x^m u_{xxxx}^2 \le C.$$

Differentiating (4.93) with respect to x, and using (A_4), (A_5), Corollaries 4.16, 4.18, 4.21, 4.23, 4.25, 4.26, 4.27, 5.3, and Lemmas 4.22, 5.2, we have

$$\int_{Q_T} x^m \theta_{xxxx}^2 \le C.$$

The proof of Lemma 5.8 is complete. □

6 Proof of Theorem 2.2.2

Let $0 < T^* < \infty$ be the maximum time of existence of strong solution (ρ, u, θ) to (2.2.1)-(2.2.3). Namely, (ρ, u, θ) is a strong solution to (2.2.1)-(2.2.3) in $\mathbb{R}^3 \times [0, T]$ for any $0 < T < T^*$, but not a strong solution in $\mathbb{R}^3 \times [0, T^*]$. We shall prove Theorem 2.2.2 by using a contradiction argument. Suppose that (2.2.5) were false, i.e.

$$M := \lim \sup_{t \nearrow T^*} (\|\rho(t)\|_{L^\infty} + \int_0^t \|\rho\theta(s)\|_{L^{\frac{12}{5}}}^4 ds) < \infty. \tag{6.1}$$

The goal is to show that under the assumption (6.1), there is a bound $C > 0$ depending only on $M, \rho_0, u_0, \theta_0, \mu, \lambda, \kappa$, and T^* such that

$$\sup_{0 \le t < T^*} \|\theta(t)\|_{L^\infty} \le C. \tag{6.2}$$

With (6.2) and (6.1), we showed in our previous paper [43] that T^* is not the maximum time, which is the desired contradiction.

Throughout the rest of the section, we denote by C a generic constant depending only on ρ_0, u_0, θ_0, T^*, M, λ, μ, κ. We denote by

$$A \lesssim B$$

if there exists a generic constant C such that $A \le CB$.

Lemma 6.1 *Under the conditions of Theorem 2.2.2 and (6.1), it holds that*

$$\sup_{0 \le t \le T} \int_{\mathbb{R}^3} \rho \le C, \text{ for any } T \in [0, T^*). \tag{6.3}$$

Proof. Integrating $(2.2.1)_1$ over $\mathbb{R}^3 \times [0, t]$, for $t < T^*$, and using the assumption $\rho_0 \in L^1$, we get (6.3). □

Lemma 6.2 *Under the conditions of Theorem 2.2.2 and (6.1), if* $3\mu > \lambda$, *it holds that*

$$\sup_{0 \le t \le T} \int_{\mathbb{R}^3} \rho |u|^4 + \int_0^T \int_{\mathbb{R}^3} |u|^2 |\nabla u|^2 \, dx \le C, \tag{6.4}$$

for any $T \in (0, T^*)$.

Proof. The detailed proof of Lemma 6.2 could be found in [43], which might be slightly modified. □

Lemma 6.3 *Under the conditions of Theorem 2.2.2 and (6.1), it holds that for any* $T \in [0, T^*)$

$$\sup_{0\le t\le T}\int_{\mathbb{R}^3}(\rho|\theta|^2+|\nabla u|^2)\,dx+\int_0^T\int_{\mathbb{R}^3}(|\nabla\theta|^2+\rho|u_t|^2)\,dxdt\le C. \tag{6.5}$$

Proof. Multiplying $(2.2.1)_2$ by u_t, and integrating by parts over $\mathbb{R}^3$, we have

$$\begin{aligned}&\int_{\mathbb{R}^3}\rho|u_t|^2+\frac{1}{2}\frac{d}{dt}\int_{\mathbb{R}^3}\left(\mu|\nabla u|^2+(\mu+\lambda)|\mathrm{div}u|^2\right)\\ =&\frac{d}{dt}\int_{\mathbb{R}^3}P\mathrm{div}u-\frac{1}{2(2\mu+\lambda)}\frac{d}{dt}\int_{\mathbb{R}^3}P^2-\frac{1}{2\mu+\lambda}\int_{\mathbb{R}^3}P_tG-\int_{\mathbb{R}^3}\rho u\cdot\nabla u\cdot u_t\\ =&\sum_{i=1}^4 VVII_i,\end{aligned} \tag{6.6}$$

where $G=(2\mu+\lambda)\mathrm{div}u-P$.

Recalling $P=\rho\theta$, we obtain from $(2.2.1)_1$ and $(2.2.1)_3$

$$P_t=-\mathrm{div}(Pu)-\rho\theta\mathrm{div}u+\mu\left(\nabla u+(\nabla u)'\right):\nabla u+\lambda\mathrm{div}u\mathrm{div}u+\Delta\theta. \tag{6.7}$$

Substituting (6.7) into $VVII_3$, and using integration by parts and Hölder inequality, we have

$$\begin{aligned}VVII_3=&-\frac{1}{2\mu+\lambda}\int_{\mathbb{R}^3}Pu\cdot\nabla G+\frac{1}{2\mu+\lambda}\int_{\mathbb{R}^3}\rho\theta\mathrm{div}uG\\ &+\frac{\mu}{2\mu+\lambda}\int_{\mathbb{R}^3}\left(\nabla u+(\nabla u)'\right):(\nabla G\otimes u)+\frac{\lambda}{2\mu+\lambda}\int_{\mathbb{R}^3}\mathrm{div}uu\cdot\nabla G\\ &+\frac{1}{2\mu+\lambda}\int_{\mathbb{R}^3}\left(\mu\Delta u+(\mu+\lambda)\nabla\mathrm{div}u\right)\cdot uG+\frac{1}{2\mu+\lambda}\int_{\mathbb{R}^3}\nabla\theta\cdot\nabla G\\ &\le C\|\rho u\theta\|_{L^2}\|\nabla G\|_{L^2}+\frac{1}{2\mu+\lambda}\int_{\mathbb{R}^3}\rho\theta\mathrm{div}uG+C\|\nabla G\|_{L^2}\big\|u|\nabla u|\big\|_{L^2}\\ &+C\|\nabla G\|_{L^2}\|\nabla\theta\|_{L^2}+\frac{1}{2\mu+\lambda}\int_{\mathbb{R}^3}\left(\mu\Delta u+(\mu+\lambda)\nabla\mathrm{div}u\right)\cdot uG.\end{aligned} \tag{6.8}$$

Substituting $(2.2.1)_2$ into (6.8), and using Sobolev inequality, (6.1) and integration by parts, we have

$$\begin{aligned}VVII_3\le&C\|\nabla G\|_{L^2}\left(\|\rho u\theta\|_{L^2}+\big\|u|\nabla u|\big\|_{L^2}+\|\nabla\theta\|_{L^2}\right)+\frac{1}{2\mu+\lambda}\int_{\mathbb{R}^3}\rho u_t\cdot uG\\ &+\frac{1}{2\mu+\lambda}\int_{\mathbb{R}^3}\rho u\cdot\nabla u\cdot uG-\frac{1}{2\mu+\lambda}\int_{\mathbb{R}^3}Pu\cdot\nabla G\\ \le&C\|\nabla G\|_{L^2}\left(\|\rho u\theta\|_{L^2}+\big\|u|\nabla u|\big\|_{L^2}+\|\nabla\theta\|_{L^2}\right)+\frac{1}{6}\int_{\mathbb{R}^3}\rho|u_t|^2\\ &+C\int_{\mathbb{R}^3}\rho|u|^2|G|^2+C\big\|u|\nabla u|\big\|_{L^2}^2\\ \le&C\|\nabla G\|_{L^2}\left(\|\rho u\theta\|_{L^2}+\big\|u|\nabla u|\big\|_{L^2}+\|\nabla\theta\|_{L^2}\right)+\frac{1}{6}\int_{\mathbb{R}^3}\rho|u_t|^2\\ &+C\int_{\mathbb{R}^3}|u|^2|\nabla u|^2+C\int_{\mathbb{R}^3}\rho|u|^2|\rho\theta|^2.\end{aligned} \tag{6.9}$$

Taking *div* on both side of $(2.2.1)_2$, we get

$$\Delta G=\operatorname{div}(\rho u_t+\rho u\cdot\nabla u). \tag{6.10}$$

By (6.10) and the standard L^2-estimates together with (6.1), we get

$$\|\nabla G\|_{L^2}\lesssim\|\rho u_t\|_{L^2}+\|\rho u\cdot\nabla u\|_{L^2}\lesssim\|\sqrt{\rho}u_t\|_{L^2}+\big\||u||\nabla u|\big\|_{L^2}. \tag{6.11}$$

Substituting (6.11) into (6.9), and using Cauchy inequality, we have

$$VVII_3\le C\|\rho u\theta\|_{L^2}^2+C\big\|u|\nabla u|\big\|_{L^2}^2+C\|\nabla\theta\|_{L^2}^2+\frac{1}{3}\int_{\mathbb{R}^3}\rho|u_t|^2. \tag{6.12}$$

For the first term of the right hand side of (6.12), using Hölder inequality, Sobolev inequality and Cauchy inequality, we have

$$\begin{aligned}\|\rho u\theta\|_{L^2}^2\le&\big\||u|^2\big\|_{L^6}\|\rho\theta\|_{L^{\frac{12}{5}}}^2\le C\big\|u|\nabla u|\big\|_{L^2}\|\rho\theta\|_{L^{\frac{12}{5}}}^2\\ \le&C\big\|u|\nabla u|\big\|_{L^2}^2+C\|\rho\theta\|_{L^{\frac{12}{5}}}^4.\end{aligned} \tag{6.13}$$

Substituting (6.13) into (6.12), we have

$$VVII_3\le C\|\rho\theta\|_{L^{\frac{12}{5}}}^4+C\big\|u|\nabla u|\big\|_{L^2}^2+C\|\nabla\theta\|_{L^2}^2+\frac{1}{3}\int_{\mathbb{R}^3}\rho|u_t|^2. \tag{6.14}$$

For $VVII_4$, using Cauchy inequality and (6.1), we have

$$VVII_4\le\frac{1}{6}\int_{\mathbb{R}^3}\rho|u_t|^2+C\int_{\mathbb{R}^3}|u|^2|\nabla u|^2. \tag{6.15}$$

Putting (6.14) and (6.15) into (6.6), and integrating it over $[0,t]$, for $t<T^*$, we have

$$\begin{aligned}&\int_0^t\int_{\mathbb{R}^3}\rho|u_t|^2+\int_{\mathbb{R}^3}\big(\mu|\nabla u|^2+(\mu+\lambda)|\operatorname{div}u|^2\big)\\ \le&2\int_{\mathbb{R}^3}P\operatorname{div}u+C\int_0^t\|\nabla\theta\|_{L^2}^2+C\\ \le&(\mu+\lambda)\int_{\mathbb{R}^3}|\operatorname{div}u|^2+C\big(\int_{\mathbb{R}^3}\rho\theta^2+\int_0^t\int_{\mathbb{R}^3}|\nabla\theta|^2\big)+C,\end{aligned}$$

where we have used Cauchy inequality, (6.1) and (6.4). Therefore,

$$\int_0^t\int_{\mathbb{R}^3}\rho|u_t|^2+\int_{\mathbb{R}^3}|\nabla u|^2\le C\big(\int_{\mathbb{R}^3}\rho\theta^2+\int_0^t\int_{\mathbb{R}^3}|\nabla\theta|^2\big)+C. \tag{6.16}$$

Multiplying $(2.2.1)_3$ by θ and integrating by parts over $\mathbb{R}^3$, we have

$$\begin{aligned}&\int_{\mathbb{R}^3}|\nabla\theta|^2+\frac{1}{2}\frac{d}{dt}\int_{\mathbb{R}^3}\rho|\theta|^2\\ =&-\int_{\mathbb{R}^3}\rho\theta^2\operatorname{div}u+\int_{\mathbb{R}^3}\frac{\mu}{2}|\nabla u+(\nabla u)'|^2\theta+\int_{\mathbb{R}^3}\lambda|\operatorname{div}u|^2\theta\\ =&\sum_{i=1}^3VVIII_i,\end{aligned} \tag{6.17}$$

For $VVIII_2$ and $VVIII_3$, we have

$$\begin{aligned}
VVIII_2+VVIII_3=&\int_{\mathbb{R}^3}\mu\left(\nabla u+(\nabla u)'\right):\nabla u\theta+\int_{\mathbb{R}^3}\lambda|\mathrm{div}u|^2\theta\\
=&-\int_{\mathbb{R}^3}\mu(\triangle u+\nabla\mathrm{div}u)\cdot u\theta-\int_{\mathbb{R}^3}\mu\left(\nabla u+(\nabla u)'\right):(\nabla\theta\otimes u)\\
&-\int_{\mathbb{R}^3}\lambda u\cdot\nabla\mathrm{div}u\theta-\int_{\mathbb{R}^3}\lambda\mathrm{div}u u\cdot\nabla\theta\\
=&-\int_{\mathbb{R}^3}(\rho u_t+\rho u\cdot\nabla u+\nabla P)\cdot u\theta-\int_{\mathbb{R}^3}\mu\left(\nabla u+(\nabla u)'\right):(\nabla\theta\otimes u)\\
&-\int_{\mathbb{R}^3}\lambda\mathrm{div}u u\cdot\nabla\theta\\
=&-\int_{\mathbb{R}^3}\rho u_t\cdot u\theta-\int_{\mathbb{R}^3}\rho(u\cdot\nabla)u\cdot u\theta+\int_{\mathbb{R}^3}\rho\theta^2\mathrm{div}u+\int_{\mathbb{R}^3}\rho\theta u\cdot\nabla\theta\\
&-\int_{\mathbb{R}^3}\mu\left(\nabla u+(\nabla u)'\right):(\nabla\theta\otimes u)-\int_{\mathbb{R}^3}\lambda\mathrm{div}u u\cdot\nabla\theta,
\end{aligned}\tag{6.18}$$

where we have used integration by parts and $(2.2.1)_2$.
Using Hölder inequality, Cauchy inequality, (6.1) and (6.4), we have

$$\begin{aligned}
&VVIII_2+VVIII_3\\
\leq&\|\sqrt{\rho}u_t\|_{L^2}\|\sqrt[4]{\rho}u\|_{L^4}\|\sqrt[4]{\rho}\theta\|_{L^4}+\big\|u|\nabla u|\big\|_{L^2}\|\rho u\|_{L^3}\|\theta\|_{L^6}+\int_{\mathbb{R}^3}\rho\theta^2\mathrm{div}u\\
&+\|\nabla\theta\|_{L^2}\|\sqrt{\rho}u\|_{L^4}\|\sqrt{\rho}\theta\|_{L^4}+C\big\|u|\nabla u|\big\|_{L^2}\|\nabla\theta\|_{L^2}\\
\leq&C\|\sqrt{\rho}u_t\|_{L^2}\|\sqrt[4]{\rho}\theta\|_{L^4}+\frac{1}{2}\|\nabla\theta\|_{L^2}^2+C\big\|u|\nabla u|\big\|_{L^2}^2+\int_{\mathbb{R}^3}\rho\theta^2\mathrm{div}u+C\|\sqrt{\rho}\theta\|_{L^4}^2.
\end{aligned}\tag{6.19}$$

Substituting (6.19) into (6.17), we have

$$\int_{\mathbb{R}^3}|\nabla\theta|^2+\frac{d}{dt}\int_{\mathbb{R}^3}\rho|\theta|^2\leq C\|\sqrt{\rho}u_t\|_{L^2}\|\sqrt[4]{\rho}\theta\|_{L^4}+C\big\|u|\nabla u|\big\|_{L^2}^2+C\|\sqrt{\rho}\theta\|_{L^4}^2.\tag{6.20}$$

Integrating (6.20) over $[0,t]$ $(t<T^*)$, and using (6.4), we have

$$\int_0^t\int_{\mathbb{R}^3}|\nabla\theta|^2+\int_{\mathbb{R}^3}\rho|\theta|^2\leq C\int_0^t\|\sqrt{\rho}u_t\|_{L^2}\|\sqrt[4]{\rho}\theta\|_{L^4}+C\int_0^t\|\sqrt{\rho}\theta\|_{L^4}^2+C.\tag{6.21}$$

Multiplying (6.21) by $2C$, and adding the resulting inequality into (6.16), we have

$$\begin{aligned}
&\int_0^t\int_{\mathbb{R}^3}\rho|u_t|^2+\int_{\mathbb{R}^3}|\nabla u|^2+\int_0^t\int_{\mathbb{R}^3}|\nabla\theta|^2+\int_{\mathbb{R}^3}\rho|\theta|^2\\
\lesssim&\int_0^t\|\sqrt{\rho}u_t\|_{L^2}\|\sqrt[4]{\rho}\theta\|_{L^4}+\int_0^t\|\sqrt{\rho}\theta\|_{L^4}^2+1\\
\leq&\frac{1}{2}\int_0^t\int_{\mathbb{R}^3}\rho|u_t|^2+C\int_0^t\|\sqrt{\rho}\theta\|_{L^2}^{\frac{1}{2}}\|\sqrt[6]{\rho}\theta\|_{L^6}^{\frac{3}{2}}+C\\
\leq&\frac{1}{2}\int_0^t\int_{\mathbb{R}^3}\rho|u_t|^2+\frac{1}{2}\int_0^t\int_{\mathbb{R}^3}|\nabla\theta|^2+C\int_0^t\int_{\mathbb{R}^3}\rho|\theta|^2+C,
\end{aligned}$$

where we have used Young inequality, Hölder inequality, Sobolev inequality and (6.1). This, together with Gronwall inequality, gives (6.5). □

Lemma 6.4 *Under the conditions of Theorem 2.2.2 and (6.1), it holds that for any* $t \in (0, T^*)$

$$\int_{\mathbb{R}^3}(|\nabla\theta|^2+\rho|\dot{u}|^2)+\int_0^t\int_{\mathbb{R}^3}(\rho|\dot{\theta}|^2+|\nabla\dot{u}|^2)\le C. \tag{6.22}$$

Proof. By [43] (see (4.35) therein), we accurately have

$$\begin{aligned}
&\frac{1}{2}\frac{d}{dt}\int_{\mathbb{R}^3}\rho|\dot{u}|^2+\int_{\mathbb{R}^3}\left(\mu|\nabla\dot{u}|^2+(\mu+\lambda)|\mathrm{div}\,\dot{u}|^2\right)\\
=&\int_{\mathbb{R}^3}(P_t\mathrm{div}\,\dot{u}+u\otimes\nabla P:\nabla\dot{u})+\mu\int_{\mathbb{R}^3}\Big(\mathrm{div}\,(\Delta u\otimes u)-\Delta(u\cdot\nabla u)\Big)\cdot\dot{u}\\
&+(\mu+\lambda)\int_{\mathbb{R}^3}\Big(\mathrm{div}\,(\nabla\mathrm{div}\,u\otimes u)-\nabla\mathrm{div}\,(u\cdot\nabla u)\Big)\cdot\dot{u}=\sum_{i=1}^3 VVIV_i.
\end{aligned}\tag{6.23}$$

For $VVIV_1$, using $(2.2.1)_1$, integration by parts, (6.1) and Hölder inequality, we have

$$\begin{aligned}
VVIV_1=&\int_{\mathbb{R}^3}\Big((\rho\theta)_t\mathrm{div}\,\dot{u}-\rho\theta(\nabla u)^t:\nabla\dot{u}-\rho\theta u\cdot\nabla\mathrm{div}\,\dot{u}\Big)\\
=&\int_{\mathbb{R}^3}\Big(\rho\dot{\theta}\mathrm{div}\,\dot{u}-\rho\theta(\nabla u)^t:\nabla\dot{u}\Big)\\
\lesssim&\|\sqrt{\rho}\dot{\theta}\|_{L^2}\|\nabla\dot{u}\|_{L^2}+\|\sqrt[4]{\rho}\theta\|_{L^4}\|\nabla u\|_{L^4}\|\nabla\dot{u}\|_{L^2}.
\end{aligned}\tag{6.24}$$

For $VVIV_2$ and $VVIV_3$, by [43] (see (4.37) and (4.38) therein), we have we have

$$VVIV_2+VVIV_3\lesssim\|\nabla\dot{u}\|_{L^2}\|\nabla u\|_{L^4}^2. \tag{6.25}$$

Substituting (6.24) and (6.25) into (6.23), and using Cauchy inequality and (6.1), we have

$$\frac{1}{2}\frac{d}{dt}\int_{\mathbb{R}^3}\rho|\dot{u}|^2+\int_{\mathbb{R}^3}\left(\mu|\nabla\dot{u}|^2+(\mu+\lambda)|\mathrm{div}\,\dot{u}|^2\right)\le\frac{\mu}{2}\|\nabla\dot{u}\|_{L^2}^2+C\|\sqrt{\rho}\dot{\theta}\|_{L^2}^2+C\|\sqrt[4]{\rho}\theta\|_{L^4}^4+C\|\nabla u\|_{L^4}^4.$$

Integrating this inequality over $[0,t]$ for $t\in(0,T^*)$, and using (6.1), (6.3), Hölder inequality and Sobolev inequality, we have

$$\int_{\mathbb{R}^3}\rho|\dot{u}|^2+\int_0^t\int_{\mathbb{R}^3}|\nabla\dot{u}|^2\le C\int_0^t\|\sqrt{\rho}\dot{\theta}\|_{L^2}^2+C\int_0^t\left(\|\nabla u\|_{L^4}^4+\|\nabla\theta\|_{L^2}^4\right). \tag{6.26}$$

The next step is to get some estimates for θ. We rewrite $(2.2.1)_3$ as follows:

$$\rho\dot{\theta}+\rho\theta\mathrm{div}u=\frac{\mu}{2}\left|\nabla u+(\nabla u)'\right|^2+\lambda(\mathrm{div}u)^2+\Delta\theta. \tag{6.27}$$

Multiplying (6.27) by $\dot{\theta}$ and integrating by parts over $\mathbb{R}^3$, we have

$$\begin{aligned}
&\int_{\mathbb{R}^3}\rho|\dot{\theta}|^2+\frac{1}{2}\frac{d}{dt}\int_{\mathbb{R}^3}|\nabla\theta|^2\\
=&-\int_{\mathbb{R}^3}\rho\theta\mathrm{div}u\dot{\theta}+\int_{\mathbb{R}^3}\Big(\frac{\mu}{2}\left|\nabla u+(\nabla u)'\right|^2+\lambda(\mathrm{div}u)^2\Big)\theta_t\\
&+\int_{\mathbb{R}^3}\Big(\frac{\mu}{2}\left|\nabla u+(\nabla u)'\right|^2+\lambda(\mathrm{div}u)^2\Big)u\cdot\nabla\theta+\int_{\mathbb{R}^3}\Delta\theta u\cdot\nabla\theta\\
=&\sum_{i=1}^4 VVV_i.
\end{aligned}\tag{6.28}$$

For VVV_1, using Hölder inequality, (6.1) and Cauchy inequality, we have

$$VVV_1 \leq C\|\sqrt{\rho}\dot{\theta}\|_{L^2}\|\nabla u\|_{L^4}\|\rho^{\frac{1}{4}}\theta\|_{L^4} \leq \frac{1}{8}\|\sqrt{\rho}\dot{\theta}\|_{L^2}^2 + C\|\nabla u\|_{L^4}^4 + C\|\nabla\theta\|_{L^2}^4. \tag{6.29}$$

For VVV_2, we have

$$\begin{aligned}
VVV_2 =& \frac{d}{dt}\int_{\mathbb{R}^3}\left(\frac{\mu}{2}\left|\nabla u + (\nabla u)'\right|^2 + \lambda(\operatorname{div}u)^2\right)\theta - \mu\int_{\mathbb{R}^3}\left(\nabla u + (\nabla u)'\right) : \left(\nabla u_t + (\nabla u_t)'\right)\theta \\
&- 2\lambda\int_{\mathbb{R}^3}\operatorname{div}u\operatorname{div}u_t\theta \\
=& \frac{d}{dt}\int_{\mathbb{R}^3}\left(\frac{\mu}{2}\left|\nabla u + (\nabla u)'\right|^2 + \lambda(\operatorname{div}u)^2\right)\theta - \mu\int_{\mathbb{R}^3}\left(\nabla u + (\nabla u)'\right) : \left(\nabla\dot{u} + (\nabla\dot{u})'\right)\theta \\
&+ \mu\int_{\mathbb{R}^3}\left(\nabla u + (\nabla u)'\right) : \left(\nabla u\cdot\nabla u + (\nabla u\cdot\nabla u)'\right)\theta \\
&+ \mu\int_{\mathbb{R}^3}\left(\nabla u + (\nabla u)'\right)\cdot(u\cdot\nabla)\left(\nabla u + (\nabla u)'\right)\theta - 2\lambda\int_{\mathbb{R}^3}\operatorname{div}u\operatorname{div}\dot{u}\theta \\
&+ 2\lambda\int_{\mathbb{R}^3}\operatorname{div}u(\nabla u)' : \nabla u\theta + 2\lambda\int_{\mathbb{R}^3}u\cdot\nabla\operatorname{div}u\operatorname{div}u\theta.
\end{aligned}$$

Using integration by parts, we have

$$\begin{aligned}
VVV_2 =& \frac{d}{dt}\int_{\mathbb{R}^3}\left(\frac{\mu}{2}\left|\nabla u + (\nabla u)'\right|^2 + \lambda(\operatorname{div}u)^2\right)\theta - \mu\int_{\mathbb{R}^3}\left(\nabla u + (\nabla u)'\right) : \left(\nabla\dot{u} + (\nabla\dot{u})'\right)\theta \\
&+ \mu\int_{\mathbb{R}^3}\left(\nabla u + (\nabla u)'\right) : \left(\nabla u\cdot\nabla u + (\nabla u\cdot\nabla u)'\right)\theta - \mu\int_{\mathbb{R}^3}\frac{|\nabla u + (\nabla u)'|^2}{2}\operatorname{div}u\theta \\
&- \mu\int_{\mathbb{R}^3}\frac{|\nabla u + (\nabla u)'|^2}{2}u\cdot\nabla\theta - 2\lambda\int_{\mathbb{R}^3}\operatorname{div}u\operatorname{div}\dot{u}\theta + 2\lambda\int_{\mathbb{R}^3}\operatorname{div}u(\nabla u)' : \nabla u\theta \\
&- \lambda\int_{\mathbb{R}^3}(\operatorname{div}u)^3\theta - \lambda\int_{\mathbb{R}^3}|\operatorname{div}u|^2u\cdot\nabla\theta \\
=& \sum_{i=1}^{9}VVV_{2,i}.
\end{aligned} \tag{6.30}$$

For $VVV_{2,2}$ and $VVV_{2,6}$, using Hölder inequality, Sobolev inequality, we have

$$VVV_{2,2} + VVV_{2,6} \lesssim \|\nabla\dot{u}\|_{L^2}\|\nabla u\|_{L^3}\|\theta\|_{L^6} \lesssim \|\nabla\dot{u}\|_{L^2}\|\nabla u\|_{L^3}\|\nabla\theta\|_{L^2}. \tag{6.31}$$

Since $\nabla u = \nabla\Delta^{-1}\left(\nabla\operatorname{div}u - \nabla\times\operatorname{curl}u\right)$, we apply Calderon-Zygmund inequality to get

$$\|\nabla u\|_{L^3} \lesssim \|\operatorname{curl}u\|_{L^3} + \|\operatorname{div}u\|_{L^3}. \tag{6.32}$$

Taking curl on both sides of $(2.2.1)_2$, we have

$$\mu\Delta(\operatorname{curl}u) = \operatorname{curl}(\rho\dot{u}). \tag{6.33}$$

By (6.33), the L^2-estimates of the elliptic equations and (6.1), we have

$$\|\nabla\operatorname{curl}u\|_{L^2} \lesssim \|\rho\dot{u}\|_{L^2} \lesssim \|\sqrt{\rho}\dot{u}\|_{L^2}. \tag{6.34}$$

By (6.10), (6.32) and (6.34), together with Sobolev inequality, we have

$$\begin{aligned}
\|\nabla u\|_{L^3} \lesssim& \|\operatorname{curl}u\|_{L^3} + \|G\|_{L^3} + \|\rho\theta\|_{L^3} \lesssim \|\operatorname{curl}u\|_{H^1} + \|G\|_{H^1} + \|\rho\|_{L^6}\|\theta\|_{L^6} \\
\lesssim& \|\operatorname{curl}u\|_{L^2} + \|\operatorname{div}u\|_{L^2} + \|\nabla(\operatorname{curl}u)\|_{L^2} + \|\nabla G\|_{L^2} + \|\nabla\theta\|_{L^2} \\
\lesssim& \|\nabla(\operatorname{curl}u)\|_{L^2} + \|\nabla G\|_{L^2} + \|\nabla\theta\|_{L^2} + 1 \\
\lesssim& \|\sqrt{\rho}\dot{u}\|_{L^2} + \|\nabla\theta\|_{L^2} + 1,
\end{aligned} \tag{6.35}$$

where we have used (6.1), (6.3) and (6.5). Substituting (6.35) into (6.31) and using Young inequality, we obtain

$$VVV_{2,2}+VVV_{2,6}\lesssim\|\nabla\dot{u}\|_{L^2}\left(\|\sqrt{\rho}\dot{u}\|_{L^2}+\|\nabla\theta\|_{L^2}+1\right)\|\nabla\theta\|_{L^2}. \tag{6.36}$$

For $VVV_{2,3}$, $VVV_{2,4}$, $VVV_{2,7}$ and $VVV_{2,8}$, using Hölder inequality, Sobolev inequality and Calderon-Zygmund inequality, we have

$$\begin{aligned}&VVV_{2,3}+VVV_{2,4}+VVV_{2,7}+VVV_{2,8}\\ \lesssim&\int_{\mathbb{R}^3}|\nabla u|^3|\theta|\lesssim\|\nabla u\|^3_{L^{\frac{18}{5}}}\|\theta\|_{L^6}\lesssim\|\nabla u\|^3_{L^{\frac{18}{5}}}\|\nabla\theta\|_{L^2}\\ \lesssim&\|\mathrm{curl}u\|^3_{L^{\frac{18}{5}}}\|\nabla\theta\|_{L^2}+\|\mathrm{div}u\|^3_{L^{\frac{18}{5}}}\|\nabla\theta\|_{L^2}\\ \lesssim&\|\mathrm{curl}u\|^3_{L^{\frac{18}{5}}}\|\nabla\theta\|_{L^2}+\|G\|^3_{L^{\frac{18}{5}}}\|\nabla\theta\|_{L^2}+\|\rho\theta\|^3_{L^{\frac{18}{5}}}\|\nabla\theta\|_{L^2}.\end{aligned} \tag{6.37}$$

Using Hölder inequality again, together with (6.1), (6.3), Sobolev inequality, Gagliardo-Nirenberg inequality, (6.5), (6.10) and (6.34), we get

$$\begin{aligned}&VVV_{2,3}+VVV_{2,4}+VVV_{2,7}+VVV_{2,8}\\ \lesssim&\|\mathrm{curl}u\|_{L^2}\|\nabla\mathrm{curl}u\|^2_{L^2}\|\nabla\theta\|_{L^2}+\|G\|_{L^2}\|\nabla G\|^2_{L^2}\|\nabla\theta\|_{L^2}+\|\rho\|^3_{L^9}\|\theta\|^3_{L^6}\|\nabla\theta\|_{L^2}\\ \lesssim&\|\sqrt{\rho}\dot{u}\|^2_{L^2}\|\nabla\theta\|_{L^2}+\|\nabla\theta\|^4_{L^2}.\end{aligned} \tag{6.38}$$

For $VVV_{2,5}$ and $VVV_{2,9}$, using Hölder inequality, Cauchy inequality, Sobolev inequality, Gagliardo-Nirenberg inequality and (6.5), we have

$$\begin{aligned}VVV_{2,5}+VVV_{2,9}\lesssim&\int_{\mathbb{R}^3}|\nabla u|^2|u||\nabla\theta|\lesssim\|\nabla u\|^2_{L^4}\|u\|_{L^6}\|\nabla\theta\|_{L^3}\\ \lesssim&\|\nabla u\|^4_{L^4}+\|\nabla u\|^2_{L^2}\|\nabla\theta\|_{L^2}\|\nabla^2\theta\|_{L^2}\\ \le&C\|\nabla u\|^4_{L^4}+C\|\nabla\theta\|_{L^2}\|\nabla^2\theta\|_{L^2}.\end{aligned} \tag{6.39}$$

From the standard elliptic estimates and (6.27), we have

$$\begin{aligned}\|\nabla^2\theta\|_{L^2}\lesssim&\|\rho\dot{\theta}\|_{L^2}+\|\rho\theta\mathrm{div}u\|_{L^2}+\|\nabla u\|^2_{L^4}\\ \lesssim&\|\sqrt{\rho}\dot{\theta}\|_{L^2}+\|\rho\|_{L^{12}}\|\theta\|_{L^6}\|\nabla u\|_{L^4}+\|\nabla u\|^2_{L^4}\\ \le&C\|\sqrt{\rho}\dot{\theta}\|_{L^2}+C\|\nabla u\|^2_{L^4}+C\|\nabla\theta\|^2_{L^2},\end{aligned} \tag{6.40}$$

where we have used Hölder inequality, (6.1), (6.3), Sobolev inequality and Cauchy inequality. Substituting (6.40) into (6.39), and using Cauchy inequality, we have

$$VVV_{2,5}+VVV_{2,9}\le\frac{1}{8}\|\sqrt{\rho}\dot{\theta}\|^2_{L^2}+C\|\nabla u\|^4_{L^4}+C\|\nabla\theta\|^4_{L^2}+C. \tag{6.41}$$

Substituting (6.36), (6.38) and (6.41) into (6.30), and using Cauchy inequality, we have

$$\begin{aligned}VVV_2\le&\frac{d}{dt}\int_{\mathbb{R}^3}\left(\frac{\mu}{2}\left|\nabla u+(\nabla u)'\right|^2+\lambda(\mathrm{div}u)^2\right)\theta\\ &+C\|\nabla\dot{u}\|_{L^2}\left(\|\sqrt{\rho}\dot{u}\|_{L^2}+\|\nabla\theta\|_{L^2}+1\right)\|\nabla\theta\|_{L^2}+C\|\sqrt{\rho}\dot{u}\|^2_{L^2}\|\nabla\theta\|_{L^2}\\ &+C\|\nabla\theta\|^4_{L^2}+\frac{1}{8}\|\sqrt{\rho}\dot{\theta}\|^2_{L^2}+C\|\nabla u\|^4_{L^4}+C.\end{aligned} \tag{6.42}$$

For VVV_3, using (6.39) and (6.41), we have

$$VVV_3\lesssim\int_{\mathbb{R}^3}|\nabla u|^2|u||\nabla\theta|\le\frac{1}{8}\|\sqrt{\rho}\dot{\theta}\|^2_{L^2}+C\|\nabla u\|^4_{L^4}+C\|\nabla\theta\|^4_{L^2}+C. \tag{6.43}$$

For VVV_4, using Hölder inequality, Sobolev inequality, Gagliardo-Nirenberg inequality, (6.5), (6.40) and Young inequality, we have

$$\begin{aligned} VVV_4 \lesssim & \|\Delta\theta\|_{L^2}\|u\|_{L^6}\|\nabla\theta\|_{L^3} \lesssim \|\Delta\theta\|_{L^2}\|\nabla u\|_{L^2}\|\nabla\theta\|_{L^2}^{\frac{1}{2}}\|\nabla^2\theta\|_{L^2}^{\frac{1}{2}} \\ \lesssim & \|\nabla\theta\|_{L^2}^{\frac{1}{2}}\|\nabla^2\theta\|_{L^2}^{\frac{3}{2}} \le \frac{1}{8}\|\sqrt{\rho}\dot{\theta}\|_{L^2}^2 + C\|\nabla\theta\|_{L^2}^4 + C\|\nabla u\|_{L^4}^4 + C. \end{aligned} \tag{6.44}$$

Putting (6.29), (6.42), (6.43) and (6.44) into (6.28), integrating the resulting inequality over $[0,t]$ for $t \in (0,T^*)$, and using Cauchy inequality, (6.1), (6.4) and (6.5), we have

$$\begin{aligned} \int_0^t\int_{\mathbb{R}^3}\rho|\dot{\theta}|^2 + \int_{\mathbb{R}^3}|\nabla\theta|^2 \le & C\int_0^t\|\nabla u\|_{L^4}^4 + C\int_{\mathbb{R}^3}|\nabla u|^2|\theta| + \varepsilon\int_0^t\|\nabla\dot{u}\|_{L^2}^2 \\ & + C_\varepsilon\int_0^t\left(\|\sqrt{\rho}\dot{u}\|_{L^2}^2 + \|\nabla\theta\|_{L^2}^2 + 1\right)\|\nabla\theta\|_{L^2}^2 + C. \end{aligned} \tag{6.45}$$

For the second term of the right hand side of (6.45), we have

$$\begin{aligned} C\int_{\mathbb{R}^3}|\nabla u|^2|\theta| \lesssim & \|\nabla u\|_{L^{\frac{12}{5}}}^2\|\theta\|_{L^6} \lesssim \|\mathrm{curl}u\|_{L^{\frac{12}{5}}}^2\|\nabla\theta\|_{L^2} + \|\mathrm{div}u\|_{L^{\frac{12}{5}}}^2\|\nabla\theta\|_{L^2} \\ \lesssim & \left(\|\mathrm{curl}u\|_{L^2}^{\frac{3}{2}}\|\nabla\mathrm{curl}u\|_{L^2}^{\frac{1}{2}} + \|G\|_{L^2}^{\frac{3}{2}}\|\nabla G\|_{L^2}^{\frac{1}{2}}\right)\|\nabla\theta\|_{L^2} + \|\rho\theta\|_{L^{\frac{12}{5}}}^2\|\nabla\theta\|_{L^2} \\ \lesssim & \|\sqrt{\rho}\dot{u}\|_{L^2}^{\frac{1}{2}}\|\nabla\theta\|_{L^2} + \|\rho\theta\|_{L^2}^{\frac{3}{2}}\|\rho\theta\|_{L^6}^{\frac{1}{2}}\|\nabla\theta\|_{L^2} \\ \lesssim & \|\sqrt{\rho}\dot{u}\|_{L^2}^{\frac{1}{2}}\|\nabla\theta\|_{L^2} + \|\nabla\theta\|_{L^2}^{\frac{3}{2}} \le \frac{1}{2}\|\nabla\theta\|_{L^2}^2 + C\|\sqrt{\rho}\dot{u}\|_{L^2} + C, \end{aligned} \tag{6.46}$$

where we have used Hölder inequality, Calderon-Zygmund inequality, Gagliardo-Nirenberg inequality, (6.1), (6.5), (6.10), (6.34), Sobolev inequality and Young inequality. Substituting (6.46) into (6.45), we have

$$\begin{aligned} \int_0^t\int_{\mathbb{R}^3}\rho|\dot{\theta}|^2 + \int_{\mathbb{R}^3}|\nabla\theta|^2 \le & C\int_0^t\|\nabla u\|_{L^4}^4 + C\|\sqrt{\rho}\dot{u}\|_{L^2} + \varepsilon\int_0^t\|\nabla\dot{u}\|_{L^2}^2 \\ & + C_\varepsilon\int_0^t\left(\|\sqrt{\rho}\dot{u}\|_{L^2}^2 + \|\nabla\theta\|_{L^2}^2 + 1\right)\|\nabla\theta\|_{L^2}^2 + C. \end{aligned} \tag{6.47}$$

Multiplying (6.47) by $2C$ and adding the resulting inequality into (6.26), we have

$$\begin{aligned} & C\int_0^t\int_{\mathbb{R}^3}\rho|\dot{\theta}|^2 + 2C\int_{\mathbb{R}^3}|\nabla\theta|^2 + \int_{\mathbb{R}^3}\rho|\dot{u}|^2 + \int_0^t\int_{\mathbb{R}^3}|\nabla\dot{u}|^2 \\ \le & 2C^2\int_0^t\|\nabla u\|_{L^4}^4 + 2C^2\|\sqrt{\rho}\dot{u}\|_{L^2} + 2\varepsilon C\int_0^t\|\nabla\dot{u}\|_{L^2}^2 \\ & + 2CC_\varepsilon\int_0^t\left(\|\sqrt{\rho}\dot{u}\|_{L^2}^2 + \|\nabla\theta\|_{L^2}^2 + 1\right)\|\nabla\theta\|_{L^2}^2 + 2C^2. \end{aligned}$$

Taking ε sufficiently small, together with Cauchy inequality, we have

$$\begin{aligned} & \int_{\mathbb{R}^3}(|\nabla\theta|^2 + \rho|\dot{u}|^2) + \int_0^t\int_{\mathbb{R}^3}(\rho|\dot{\theta}|^2 + |\nabla\dot{u}|^2) \\ \lesssim & \int_0^t\|\nabla u\|_{L^4}^4 + \int_0^t\left(\|\sqrt{\rho}\dot{u}\|_{L^2}^2 + \|\nabla\theta\|_{L^2}^2 + 1\right)\|\nabla\theta\|_{L^2}^2 + 1. \end{aligned} \tag{6.48}$$

For the first term of the right hand side of (6.48), similar to (6.35), we have

$$\begin{aligned}\int_0^t \|\nabla u\|_{L^4}^4 &\lesssim \int_0^t \|\mathrm{curl}u\|_{L^4}^4 + \int_0^t \|G\|_{L^4}^4 + \int_0^t \|\nabla\theta\|_{L^2}^4 \\ &\lesssim \int_0^t \|\mathrm{curl}u\|_{L^2}\|\nabla\mathrm{curl}u\|_{L^2}^3 + \int_0^t \|G\|_{L^2}\|\nabla G\|_{L^2}^3 + \int_0^t \|\nabla\theta\|_{L^2}^4 \\ &\lesssim \int_0^t \|\sqrt{\rho}\dot{u}\|_{L^2}^3 + \int_0^t \|\nabla\theta\|_{L^2}^4.\end{aligned} \tag{6.49}$$

By (6.48), (6.49) and Cauchy inequality, we have

$$\begin{aligned}&\int_{\mathbb{R}^3}(|\nabla\theta|^2+\rho|\dot{u}|^2) + \int_0^t\int_{\mathbb{R}^3}(\rho|\dot{\theta}|^2+|\nabla\dot{u}|^2) \\ \lesssim &\int_0^t \|\sqrt{\rho}\dot{u}\|_{L^2}^3 + \int_0^t \left(\|\sqrt{\rho}\dot{u}\|_{L^2}^2 + \|\nabla\theta\|_{L^2}^2 + 1\right)\|\nabla\theta\|_{L^2}^2 + 1 \\ \lesssim &\int_0^t \left(\|\sqrt{\rho}\dot{u}\|_{L^2}^2 + \|\nabla\theta\|_{L^2}^2 + 1\right)\left(\|\nabla\theta\|_{L^2}^2 + \|\sqrt{\rho}\dot{u}\|_{L^2}^2\right) + 1.\end{aligned} \tag{6.50}$$

By (6.1), (6.4), (6.5), we have

$$\int_0^t \left(\|\sqrt{\rho}\dot{u}\|_{L^2}^2 + \|\nabla\theta\|_{L^2}^2\right) \le C,$$

for any $t \in (0, T^*)$. This, together with (6.50) and Gronwall inequality, deduces (6.22). □

Corollary 6.5 *Under the conditions of Theorem 2.2.2 and (6.1), it holds that for any $T \in (0, T^*)$*

$$\sup_{0\le t\le T}(\|\nabla G\|_{L^2} + \|\nabla\mathrm{curl}u\|_{L^2} + \|\nabla u\|_{L^6} + \|u\|_{L^\infty}) + \int_0^T\int_{\mathbb{R}^3}|\nabla^2\theta|^2 \le C. \tag{6.51}$$

Proof. The detailed proof of the lemma could be found in [43] (see Corollary 4.5 therein). □

Lemma 6.6 *Under the conditions of Theorem 2.2.2 and (6.1), it holds that for any $t \in (0, T^*)$*

$$\int_{\mathbb{R}^3}\rho|\theta_t|^2 + \int_0^t\int_{\mathbb{R}^3}|\nabla\theta_t|^2 \le C. \tag{6.52}$$

Proof. By [43], we have

$$\begin{aligned}&\frac{1}{2}\frac{d}{dt}\int_{\mathbb{R}^3}\rho|\theta_t|^2 + \int_{\mathbb{R}^3}|\nabla\theta_t|^2 \\ =&-\int_{\mathbb{R}^3}\rho_t\left(\frac{\theta_t}{2} + u\cdot\nabla\theta + \theta\mathrm{div}u\right)\theta_t - \int_{\mathbb{R}^3}\rho(u_t\cdot\nabla\theta + u\cdot\nabla\theta_t + \theta_t\mathrm{div}u)\theta_t \\ &-\int_{\mathbb{R}^3}\rho\theta\mathrm{div}u_t\theta_t + \mu\int_{\mathbb{R}^3}\left(\nabla u + (\nabla u)'\right):\left(\nabla u_t + (\nabla u_t)'\right)\theta_t + 2\lambda\int_{\mathbb{R}^3}\mathrm{div}u\mathrm{div}u_t\theta_t \\ =&\sum_{i=1}^5 VVVI_i,\end{aligned} \tag{6.53}$$

and

$$\begin{aligned}
VVVI_1 =& -\int_{\mathbb{R}^3} \rho u\cdot\nabla\theta_t\left(\frac{\theta_t}{2}+u\cdot\nabla\theta+\theta\mathrm{div}u\right)-\int_{\mathbb{R}^3}\rho u\cdot\frac{\nabla\theta_t}{2}\theta_t\\
&-\int_{\mathbb{R}^3}\rho u\cdot\left(\nabla(u\cdot\nabla)\theta+u\cdot\nabla\nabla\theta\right)\theta_t-\int_{\mathbb{R}^3}\rho u\cdot\left(\nabla\theta\mathrm{div}u+\theta\nabla\mathrm{div}u\right)\theta_t\\
=&\sum_{i=1}^{4}VVVI_{1,i}.
\end{aligned}\tag{6.54}$$

For $VVVI_{1,1}$, we have

$$\begin{aligned}
VVVI_{1,1}\le&\frac{1}{24}\|\nabla\theta_t\|_{L^2}^2+C\|\sqrt{\rho}\theta_t\|_{L^2}^2+C\|\nabla\theta\|_{L^2}^2+C\|\sqrt[4]{\rho}\theta\|_{L^4}^2\|\nabla u\|_{L^4}^2\\
\le&\frac{1}{24}\int_{\mathbb{R}^3}|\nabla\theta_t|^2+C\int_{\mathbb{R}^3}\rho|\theta_t|^2+C,
\end{aligned}\tag{6.55}$$

where we have used Cauchy inequality, (6.1), (6.5), (6.22) and (6.51).
For $VVVI_{1,2}$, using Cauchy inequality, (6.1) and (6.51) again, we have

$$VVVI_{1,2}\le\frac{1}{24}\int_{\mathbb{R}^3}|\nabla\theta_t|^2+C\int_{\mathbb{R}^3}\rho|\theta_t|^2.\tag{6.56}$$

For $VVVI_{1,3}$, by [43] (see (4.65) therein), we have

$$VVVI_{1,3}\lesssim\int_{\mathbb{R}^3}\rho|\theta_t|^2+\int_{\mathbb{R}^3}|\nabla^2\theta|^2+1.\tag{6.57}$$

For $VVVI_{1,4}$, integrating by parts, we have

$$\begin{aligned}
VVVI_{1,4}=&-\int_{\mathbb{R}^3}\rho u\cdot\nabla\theta\mathrm{div}u\theta_t-\frac{1}{2\mu+\lambda}\int_{\mathbb{R}^3}\rho\theta u\cdot\nabla G\theta_t-\frac{1}{2\mu+\lambda}\int_{\mathbb{R}^3}\rho\theta u\cdot\nabla(\rho\theta)\theta_t\\
=&-\int_{\mathbb{R}^3}\rho u\cdot\nabla\theta\mathrm{div}u\theta_t-\frac{1}{2\mu+\lambda}\int_{\mathbb{R}^3}\rho\theta u\cdot\nabla G\theta_t\\
&+\frac{1}{2(2\mu+\lambda)}\int_{\mathbb{R}^3}\rho^2\theta^2u\cdot\nabla\theta_t+\frac{1}{2(2\mu+\lambda)}\int_{\mathbb{R}^3}\rho^2\theta^2\mathrm{div}u\theta_t.
\end{aligned}\tag{6.58}$$

Furthermore, we can get

$$\begin{aligned}
VVVI_{1,4}\lesssim&\|\sqrt{\rho}\theta_t\|_{L^2}\|\nabla u\|_{L^6}\|\nabla\theta\|_{L^3}+\|\nabla G\|_{L^2}\|\theta_t\|_{L^6}\|\theta\|_{L^6}\|\rho\|_{L^6}\\
&+\|\theta\|_{L^6}^2\|\rho\|_{L^{12}}^2\|\nabla\theta_t\|_{L^2}+\|\nabla u\|_{L^3}\|\theta_t\|_{L^6}\|\theta\|_{L^6}^2\|\rho\|_{L^{12}}^2\\
\le&\frac{1}{24}\|\nabla\theta_t\|_{L^2}^2+C\|\nabla^2\theta\|_{L^2}^2+C\|\sqrt{\rho}\theta_t\|_{L^2}^2+C,
\end{aligned}\tag{6.59}$$

where we have used Hölder inequality, Sobolev inequality, (6.1), (6.3), (6.5), (6.22), (6.51) and Cauchy inequality. Substituting (6.55), (6.56), (6.57) and (6.59) into (6.54), we have

$$VVVI_1\le\frac{1}{8}\int_{\mathbb{R}^3}|\nabla\theta_t|^2+C\int_{\mathbb{R}^3}\rho|\theta_t|^2+C\int_{\mathbb{R}^3}|\nabla^2\theta|^2+C.\tag{6.60}$$

For $VVVI_2$,

$$\begin{aligned}
VVVI_2=&-\int_{\mathbb{R}^3}\rho\dot{u}\cdot\nabla\theta\theta_t+\int_{\mathbb{R}^3}\rho(u\cdot\nabla)u\cdot\nabla\theta\theta_t-\int_{\mathbb{R}^3}\rho u\cdot\nabla\theta_t\theta_t-\int_{\mathbb{R}^3}\rho|\theta_t|^2\mathrm{div}u\\
\lesssim&\|\sqrt{\rho}\dot{u}\|_{L^2}\|\theta_t\|_{L^6}\|\nabla\theta\|_{L^3}+\|\sqrt{\rho}\theta_t\|_{L^2}\|\nabla u\|_{L^6}\|\nabla\theta\|_{L^3}+\|\sqrt{\rho}\theta_t\|_{L^2}\|\nabla\theta_t\|_{L^2}\\
&+\|\sqrt{\rho}\theta_t\|_{L^2}\|\theta_t\|_{L^6}\|\nabla u\|_{L^3}\\
\le&\frac{1}{8}\int_{\mathbb{R}^3}|\nabla\theta_t|^2+C\int_{\mathbb{R}^3}\rho|\theta_t|^2+C\int_{\mathbb{R}^3}|\nabla^2\theta|^2+C,
\end{aligned}\tag{6.61}$$

where we have used Hölder inequality, Sobolev inequality, (6.1), (6.5), (6.22) and (6.51). For $VVVI_3$, integrating by parts, we have

$$\begin{aligned}VVVI_3 =& -\int_{\mathbb{R}^3} \rho\theta \mathrm{div}\dot{u}\theta_t + \int_{\mathbb{R}^3} \rho\theta \mathrm{div}(u\cdot\nabla u)\theta_t\\ =& -\int_{\mathbb{R}^3} \rho\theta \mathrm{div}\dot{u}\theta_t + \int_{\mathbb{R}^3} \rho\theta\nabla u:(\nabla u)'\theta_t + \int_{\mathbb{R}^3} \rho\theta u\cdot\nabla\mathrm{div}u\theta_t\\ =& -\int_{\mathbb{R}^3} \rho\theta \mathrm{div}\dot{u}\theta_t + \int_{\mathbb{R}^3} \rho\theta\nabla u:(\nabla u)'\theta_t + \frac{1}{2\mu+\lambda}\int_{\mathbb{R}^3} \rho\theta u\cdot\nabla G\theta_t\\ &- \frac{1}{2\mu+\lambda}\int_{\mathbb{R}^3}\frac{\rho^2}{2}\theta^2\mathrm{div}u\theta_t - \frac{1}{2\mu+\lambda}\int_{\mathbb{R}^3}\frac{\rho^2}{2}\theta^2 u\cdot\nabla\theta_t.\end{aligned} \tag{6.62}$$

Furthermore, using Hölder inequality, Sobolev inequality, (6.1), (6.3), (6.5), (6.22), (6.51) and Young inequality, we have

$$\begin{aligned}VVVI_3 \lesssim& \|\nabla\dot{u}\|_{L^2}\|\theta_t\|_{L^6}\|\theta\|_{L^6}\|\rho\|_{L^6} + \|\sqrt{\rho}\theta\|_{L^2}\|\nabla u\|_{L^6}^2\|\theta_t\|_{L^6}\\ &+ \|\nabla G\|_{L^2}\|\theta_t\|_{L^6}\|\theta\|_{L^6}\|\rho\|_{L^6} + \|\nabla u\|_{L^3}\|\theta_t\|_{L^6}\|\theta\|_{L^6}^2\|\rho\|_{L^{12}}^2\\ &+ \|\theta\|_{L^6}^2\|\rho\|_{L^{12}}^2\|\nabla\theta_t\|_{L^2}\\ \le& \frac{1}{8}\|\nabla\theta_t\|_{L^2}^2 + C\|\nabla\dot{u}\|_{L^2}^2 + C.\end{aligned} \tag{6.63}$$

Similar to VVV_2, for $VVVI_4$ and $VVVI_5$, we deduce

$$\begin{aligned}VVVI_4 + VVVI_5 \le& C\|\nabla\dot{u}\|_{L^2}\|\nabla u\|_{L^3}\|\theta_t\|_{L^6} + C\int_{\mathbb{R}^3}|\nabla u|^3|\theta_t| + C\int_{\mathbb{R}^3}|\nabla u|^4 + \frac{1}{16}\int_{\mathbb{R}^3}|\nabla\theta_t|^2\\ \le& C\|\nabla\dot{u}\|_{L^2}\|\nabla\theta_t\|_{L^2} + C\|\nabla u\|_{L^{\frac{18}{5}}}^3\|\nabla\theta_t\|_{L^2} + \frac{1}{16}\int_{\mathbb{R}^3}|\nabla\theta_t|^2 + C\\ \le& \frac{1}{8}\int_{\mathbb{R}^3}|\nabla\theta_t|^2 + C\int_{\mathbb{R}^3}|\nabla\dot{u}|^2 + C,\end{aligned} \tag{6.64}$$

where we have used Hölder inequality, integration by parts, Cauchy inequality, (6.5), (6.51), the interpolation inequality and Sobolev inequality.
Putting (6.60), (6.61), (6.63) and (6.64) into (6.53), we have

$$\frac{d}{dt}\int_{\mathbb{R}^3}\rho|\theta_t|^2 + \int_{\mathbb{R}^3}|\nabla\theta_t|^2 \le C\int_{\mathbb{R}^3}\rho|\theta_t|^2 + C\int_{\mathbb{R}^3}(|\nabla\dot{u}|^2 + |\nabla^2\theta|^2) + C. \tag{6.65}$$

By (6.65), (6.22), (6.51) and Gronwall inequality, we complete the proof of Lemma 6.6. □

Corollary 6.7 *Under the conditions of Theorem 2.2.2 and (6.1), it holds that for any* $t \in (0, T^*)$

$$\int_{\mathbb{R}^3}|\nabla^2\theta|^2 \le C. \tag{6.66}$$

Proof. It follows from $(2.2.1)_3$, (6.1), (6.3), (6.5), (6.22), (6.51), (6.52) and the interpolation inequality that

$$\begin{aligned}\|\nabla^2\theta\|_{L^2} \lesssim& \|\rho\theta_t\|_{L^2} + \|\rho u\cdot\nabla\theta\|_{L^2} + \|\rho\theta\mathrm{div}u\|_{L^2} + \|\nabla u\|_{L^4}^2\\ \lesssim& \|\sqrt{\rho}\theta_t\|_{L^2} + \|\nabla\theta\|_{L^2} + \|\rho\|_{L^6}\|\theta\|_{L^6}\|\mathrm{div}u\|_{L^6} + 1\\ \le& C.\end{aligned}$$

□

By (6.22), (6.66) and Sobolev inequality, we get the following corollary which is the desired one, i.e., (6.2).

Corollary 6.8 *Under the conditions of Theorem 2.2.2 and (6.1), it holds that for any* $t \in (0, T^*)$

$$\|\theta\|_{L^\infty(0,t;L^\infty)} \leq C. \tag{6.67}$$

Acknowledgements

The research was supported by the National Natural Science Foundation of China #10625105, #11071093, the PhD specialized grant of the Ministry of Education of China #20100144110001, and the Special Fund for Basic Scientific Research of Central Colleges #CCNU10C01001.

References

[1] D. Bresch, B. Desjardins, *On the existence of global weak solutions to the Navier-Stokes equations for viscous compressible and heat conducting fluids,* J. Math. Pures Appl., 87(2007), 57-90.

[2] Y. Cho, H.J. Choe, H. Kim, *Unique solvability of the initial boundary value problems for compressible viscous fluids,* J. Math. Pures Appl., 83(2004), 243-275.

[3] Y. Cho, H. Kim, *On classical solutions of the compressible Navier-Stokes equations with nonnegative initial densities,* Manuscripta Math., 120(2006), 91-129.

[4] Y. Cho, H. Kim, *Existence results for viscous polytropic fluids with vacuum,* J. Differential Equations, 228(2006), 377-411.

[5] H.J. Choe, H. Kim, *Strong solutions of the Navier-Stokes equations for isentropic compressible fluids,* J. Differential Equations, 190(2003), 504-523.

[6] H.J. Choe, H. Kim, *Global existence of the radially symmetric solutions of the Navier-Stokes equations for the isentropic compressible fluids,* Math. Methods Appl. Sci., 28(2005), 1-28.

[7] S.J. Ding, H.Y. Wen, L. Yao, C.J. Zhu, *Global classical spherically symmetric solution of compressible isentropic Navier-Stokes equations with vacuum,* SIAM J. Math. Anal., 2011, to appear.

[8] S.J. Ding, H.Y. Wen, C.J. Zhu, *Global classical large solutions to 1D compressible Navier-Stokes equations with density-dependent viscosity and vacuum,* J. Differential Equations, 251(2011), 1696-1725.

[9] L.C. Evans, *Partial Differential Equations,* Graduate Studies in Mathematics, Amer. Math. Soc., Providence, Rhode Island, Vol. 19, 1998.

[10] J. Fan, S. Jiang, G. Ni, *Uniform boundedness of the radially symmetric solutions of the Navier-Stokes equations for isentropic compressible fluids,* Osaka J. Math., 46(2009), 863-876.

[11] J. Fan, S. Jiang, Y. Ou, *A blow-up criterion for compressible viscous heat-conductive flows,* Ann. I. H. Poincaré-AN, 27(2010), 337-350.

[12] D.Y. Fang, R.Z. Zi, T. Zhang, *A blow-up criterion for two dimensional compressible viscous heat-conductive flows.* arXiv:1107.4663v1 [math.AP] 23 Jul 2011.

[13] E. Feireisl, *On the motion of a viscous, compressible and heat conducting fluid,* Indiana Univ. Math. J., 53(2004), 1705-1738.

[14] E. Feireisl, *Dynamics of Viscous Compressible Fluids,* Oxford Univ. Press, Oxford, 2004.

[15] E. Feireisl, A. Novotný, H. Petzeltová, *On the existence of globally defined weak solutions to the Navier-Stokes equations,* J. Math. Fluid Mech., 3(2001), 358-392.

[16] H. Frid, V. Shelukhin, *Vanishing shear viscosity in the equations of compressible fluids for the flows with the cylinder symmetry,* SIAM J. Math. Anal., 31(2000), 1144-1156.

[17] Z.H. Guo, C.J. Zhu, *Global weak solutions and asymptotic behavior to 1D compressible Navier-Stokes equations with density-dependent viscosity and vacuum,* J. Differential Equations, 248(2010), 2768-2799.

[18] D. Hoff, H. Jenssen, *Symmetric nonbarotropic flows with large data and forces,* Arch. Rational Mech. Anal., 173 (2004), 297-343.

[19] X. Huang, J. Li, *Global classical and weak solutions to the three-dimensional full compressible Navier-Stokes system with vacuum and large oscillations,* arXiv:1107.4655v2 [math-ph] 26 Jul 2011.

[20] X. Huang, J. Li, Z. Xin, *Global well-posedness of classical solutions with large oscillations and vacuum to the three-dimensional isentropic compressible Navier-Stokes equations,* arXiv:1004.4749v2 [math-ph] 8 Jul 2010.

[21] N. Itaya, *On the Cauchy problem for the system of fundamental equations describing the movement of compressible viscous fluid,* Kodai Math. Sem. Rep., 23(1971), 60-120.

[22] S. Jiang, *Global spherically symmetric solutions to the equations of a viscous polytropic ideal gas in an exterior domain,* Comm. Math. Phys., 178(1996), 339-374.

[23] S. Jiang, *Global smooth solutions of the equations of a viscous, heat-conducting one-dimensional gas with density-dependent viscosity,* Math. Nachr., 190(1998), 169-183.

[24] S. Jiang, *Large-time behavior of solutions to the equations of a viscous polytropic ideal gas,* Annali di Matematica pura ed applicata, (IV)(1998), 253-275.

[25] S. Jiang, *Large-time behavior of solutions to the equations of a one-dimensional viscous polytropic ideal gas in unbounded domains,* Comm. Math. Phys., 200(1999), 181-193.

[26] S. Jiang, J.W. Zhang, *Boundary layers for the Navier-Stokes equations of compressible heat-conducting flows with cylindrical symmetry,* SIAM J. Math. Anal., 41(2009), 237-268.

[27] S. Jiang, P. Zhang, *On spherically symmetric solutions of the compressible isentropic Navier-Stokes equations,* Comm. Math. Phys., 215(2001), 559-581.

[28] S. Jiang, P. Zhang, *Global weak solutions to the Navier-Stokes equations for a 1D viscous polytropic ideal gas,* Quart. Appl. Math., 61(2003), 435-449.

[29] B. Kawohl, *Global existence of large solutions to initial boundary value problems for a viscous, heat-conducting, one-dimensional real gas,* J. Differential Equations, 58(1985), 76-103.

[30] A.V. Kazhikhov, V.V. Shelukhi, *Unique global solution with respect to time of initial-boundary value problems for one-dimensional equations of a viscous gas,* Prikl. Mat. Meh., 41(1977), 282-291.

[31] P.L. Lions, Mathematical Topics in Fluid Mechanics, Vol. II, Compressible Models, Clarendon Press, Oxford, 1998.

[32] T.P. Liu, Z. Xin, T. Yang, *Vacuum states of compressible flow,* Discrete Contin. Dyn. Syst., 4(1998), 1-32.

[33] T. Luo, Z.P. Xin, T. Yang, *Interface behavior of compressible Navier-Stokes equations with vacuum,* SIAM J. Math. Anal., 31(2000), 1175-1191.

[34] A. Matsumura, T. Nishida, *The initial value problem for the equations of motion of viscous and heat-conductive gases,* J. Math. Kyoto Univ., 20(1980), 67-104.

[35] A. Matsumura, T. Nishida, *The initial boundary value problems for the equations of motion of compressible and heat-conductive fluids,* Comm. Math. Phys., 89(1983), 445-464.

[36] X. Qin, Z. Yao, *Global smooth solutions of the compressible Navier-Stokes equations with density-dependent viscosity,* J. Differential Equations, 244(2008), 2041-2061.

[37] R. Salvi, I. Straškraba, *Global existence for viscous compressible fluids and their behavior as $t \to \infty$,* J. Fac. Sci. Univ. Tokyo, Sect. IA Math., 40(1993), 17-51.

[38] J. Simon, *Nonhomogeneous viscous incompressible fluids: existence of vecocity, density and pressure*, SIAM J. Math. Anal., 21(1990), 1093-1117.

[39] Y.Z. Sun, C. Wang, Z.F. Zhang, *A Beale-Kato-Majda criterion for three dimensional compressible viscous heat-conductive Flows.* Arch. Rational Mech. Anal., 201(2011), 727-742.

[40] A. Tani, *On the first initial-boundary value problem of compressible viscous fluid motion,* Publ. Res. Inst. Math. Sci. Kyoto Univ., 13(1977), 193-253.

[41] S.W. Vong, T. Yang, C.J. Zhu, *Compressible Navier-Stokes equations with degenerate viscosity coefficient and vacuum (II),* J. Differential Equations, 192(2003), 475-501.

[42] H.Y. Wen, C.J. Zhu, *Global classical large solutions to Navier-Stokes equations for viscous compressible and heat conducting fluids with vacuum,* arXiv:1103.1421v1 [math.AP] 8 Mar 2011, 1-38.

[43] H.Y. Wen, C.J. Zhu. *Blow-up criterions of strong solutions to 3D compressible Navier-Stokes equations with vacuum*, arXiv:1111.2657v1 [math.AP] 11 Nov 2011.

[44] Z. Xin, *Blowup of smooth solutions to the compressible Navier-Stokes equation with compact density,* Comm. Pure Appl. Math., 51(1998), 229-240.

[45] T. Yang, *Singular behavior of vacuum states for compressible fluids,* Journal of Computational and Applied Mathematics, 190(2006), 211-231.

[46] T. Yang, Z.A. Yao, C.J. Zhu, *Compressible Navier-Stokes equations with density-dependent viscosity and vacuum,* Comm. Partial Differential Equations, 26(2001), 965-981.

[47] T. Yang, C.J. Zhu, *Compressible Navier-Stokes equations with degenerate viscosity coefficient and vacuum,* Comm. Math. Phys., 230(2002), 329-363.

[48] C.J. Zhu, *Asymptotic behavior of compressible Navier-Stokes equations with density-dependent viscosity and vacuum,* Comm. Math. Phys., 293(2010), 279-299.